

SAIP2014



Contribution ID : 13

Constraining Beyond the Standard Model physics with the newly discovered Higgs boson with the ATLAS detector.

Tuesday 08 Jul 2014 at 15:20 (00h20')

Abstract :

The discovery of the Higgs boson opens many perspectives to explore physics beyond the Standard Model. This talk describes the measurements and searches performed by the ATLAS experiment using the Higgs boson as a portal to search or constrain new physics. A selected list of the presented searches include: Constraints on the existence of weakly interacting dark-matter particles through Higgs boson to invisible decays; upper limits of Flavour-Changing-Neutral-Currents in top quark decays and exclusion limits on the existence of Two Higgs Doublet Models. No significant signs of new physics were found in the data. Limits, in some cases the world's most sensitive, are placed on the different searches.

Award :

No

Level :

N/A

Supervisor :

Dr. Trevor Vickey (Trevor.Vickey@wits.ac.za) University of the Witwatersrand

Paper :

Yes

Primary authors : CARRILLO-MONTOYA, German David (University of the Witwatersrand)

Co-authors :

Presenter : CARRILLO-MONTOYA, German David (University of the Witwatersrand)

Session classification : NPRP

Track classification : Track B - Nuclear, Particle and Radiation Physics

Type : Oral Presentation