

SAIP 2011



Contribution ID : 197

Decomposition of the field within an apertured plano-concave resonator

Wednesday 13 Jul 2011 at 11:15 (00h15')

Content :

A method of experimentally decomposing a field within an apertured plano-concave resonator cavity into a forward propagating field (plane to concave) and a backward propagating field (concave to plane) is presented. The use of an intra-cavity aperture which is a simple method of laser beam shaping is a means of mode selection as higher-order modes are actively discriminated. This apertured cavity is modelled by two fundamental resonator theories, namely, Fox-Li analysis and Laguerre-Gaussian decomposition where they are used in the determination of the respective beam profiles at a specific plane. These theories traditionally differ from one another for an apertured cavity where the Gaussian beam within the cavity is truncated. A preliminary set-up is characterized for Gaussian propagation in an attempt to verify that the cavity is viable and accurate. A comparison of experimental data with the theories is presented.

Level (Hons, MSc, PhD, other)? :

MSc

Consider for a student award (Yes / No)? :

Yes

Short Paper :

No

Primary authors : Mr. NAIDOO, Darryl (Council for Scientific and Industrial Research)

Co-authors : Prof. FORBES, Andrew (Council for Scientific and Industrial Research) ; Prof. AÏT-AMEUR, Kamel (CIMAP—ENSICAEN)

Presenter : Mr. NAIDOO, Darryl (Council for Scientific and Industrial Research)

Session classification : LOS

Track classification : Track C - Lasers, Optics and Spectroscopy

Type : Oral Presentation