

SAIP2013



Contribution ID : 458

Spin-lattice relaxation in Fe implanted ZnO

Wednesday 10 Jul 2013 at 10:50 (00h20')

Abstract :

ZnO doped with 3d metals has attracted much attention since the theoretical prediction of Curie temperatures above room temperature resulting from carrier mediated magnetic interactions due to itinerant holes coupling with localized dopant spins. Several published reports reveal that these materials exhibit different forms of magnetism, the origin of which is still under debate. We have undertaken emission ^{57}Fe Mössbauer spectroscopy measurements following implantation of radioactive $^{57}\text{Mn}^+$ ions ($3 \cdot 10^{12}/\text{cm}^2$) at ISOLDE/CERN, on virgin and ^{56}Fe pre-implanted ZnO samples with fluences of $2 \cdot 10^{13}/\text{cm}^2$, $5 \cdot 10^{13}/\text{cm}^2$ and $7 \cdot 10^{13}/\text{cm}^2$. The samples were held at temperatures between 77 – 726 K in an implantation chamber and spectra were collected at gamma emission angles of 60° relative to the sample *c*-axis. The spectra reveal magnetic structure in the ‘wings’ on a wide velocity scale (± 12 mm/s) which were analysed using a semi-empirical relaxation model utilizing five Blume-Tjon sextets. The observed temperature-dependent magnetic effect may be explained by a slow spin-lattice relaxation due to paramagnetic Fe^{3+} weakly coupled to the lattice. The central part of spectra were fitted with two quadrupole split doublets assigned to Fe in regular crystalline sites (Fe_C) and in interstitial sites (Fe_I) in the lattice. The spin-lattice relaxation rates of Fe^{3+} in the ^{56}Fe pre-implanted ZnO samples increases with increasing fluence, and in the virgin ZnO sample follows a T^9 temperature dependence but in the pre-implanted samples show a transition towards a T^2 dependence. This result suggests that an increase in implantation fluence favours the two phonon process. The annealing behaviour and variation of hyperfine parameters in these materials will also be discussed.

Award :

Yes

Level :

PhD

Supervisor :

Dr Deena Naidoo Deena.Naidoo@wits.ac.za School of Physics, University of the Witwatersrand

Paper :

No

Primary authors : Mr. MASENDA, Hilary (Wits)

Co-authors : Dr. GEBURT, Sebastian (Institute of Solid State Physics, University of Jena) ; Prof. BHARUTH-RAM, Krish (DUT) ; Dr. NAIDOO, Deena (Wits) ; Prof. GUNNLAUGSSON, Haraldur Páll (Department of Physics and Astronomy, Aarhus University) ; Dr. MANTOVAN, Roberto (Laboratorio MDM, IMM-CNR,) ; Dr. JOHNSTON, Karl (PH Dept, ISOLDE/CERN) ; Dr. MØLHOLT, Torben (Science Institute, University of Iceland) ; Mr. NCUBE, Mehluli (Wits) ; Prof. RONNING, Carsten (Institute of Solid State Physics, University of Jena) ; Prof. GÍSLASON, Hafliði Pétur (Science Institute, University of Iceland) ; Prof. ÓLAFSSON, Sveinn (Science Institute, University of Iceland) ; Prof. LANGOUCHE, Guido (Instituut voor Kern-en Stralings fysika, University of Leuven) ; Prof. WEYER, Gerd (Department of Physics and Astronomy, Aarhus University)

Presenter : Mr. MASENDA, Hilary (Wits)

Session classification : DCMPM2

Track classification : Track A - Division for Condensed Matter Physics and Materials

Type : Oral Presentation