

SAIP2012



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Contribution ID : 476

A transmitter-receiver model of the propagation of the second moments of a non-Gaussian non-spherical laser beam through an ABCD system.

Wednesday 11 Jul 2012 at 11:35 (00h20')

Abstract :

A free-propagating laser beam can be completely defined if its beam size, divergence and space-angular moment at a plane are known. I present a model which is based on the idea that under these circumstances, once these parameters are known in one plane, say $z = 0$, we can calculate the same parameters at a subsequent plane, $z > 0$ if the space between the two planes is described an ABCD matrix. I show that we can calculate the beam quality factor (M^2), a parameter which does not depend on the ABCD elements. The resulting expression for M^2 is based on the so-called, curvature removal method. I also show that, during translation, the so-called local parameters (which depend on z), i.e., the divergence, curvature and the Guoy shift, and what I am going to call the global parameters (which are independent of z), i.e., the waist position, waist size, Rayleigh range and far-field divergence, can all be calculated from the input parameters if the ABCD matrix elements are known. I then derive a few well-known expressions to confirm the model's validity.

Award :

No

Paper :

No

Primary authors : Dr. MAFUSIRE, Cosmas (CSIR National Laser Centre)

Co-authors :

Presenter : Dr. MAFUSIRE, Cosmas (CSIR National Laser Centre)

Session classification : Photonics

Track classification : Track C - Photonics

Type : Oral Presentation