

SAIP2012



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Contribution ID : 65

Application of the Isobaric Mass Multiplet Equation to the rp process in Nuclear Astrophysics

Friday 13 Jul 2012 at 08:20 (00h20')

Abstract :

In many cases levels of proton-rich nuclei participating in rp processes have not been measured and one has to rely on theory to estimate the reaction rates. However, the Isobaric Mass Multiplet Equation (IMME) affords a more reliable method of obtaining levels in the final $T=1$ nucleus of a (p, γ) reaction in terms of experimental energies of the isobaric analog partners and a small coefficient c (typically about 200 keV) that can be calculated [1]. The power of the IMME method, which is mainly empirically based with a small theoretical component, to estimate energies of nuclei participating in the rp process is emphasized by way of illustrating favorable cases where the method works particularly well. In the case of ^{26}Si it is demonstrated that the theoretical component can be calculated with considerable accuracy, and in the case of ^{30}S it is shown that a large number of states can be predicted when only a small number have been experimentally measured. However, because of the semi-empirical nature of the method, it should be stressed that its application depends on knowing energies of the analog states adjacent to the nucleus in question. Alternatively, one has to resort to shell-model calculations with reliable two-body interactions. As a further illustration the method is applied to the $^{35}\text{Ar}(p,\gamma)^{36}\text{K}$ reaction.[1] W. A. Richter, B. A. Brown, A. Signoracci and M. Wiescher, Phys. Rev.C 83, 065803 (2011)

Award :

No

Paper :

No

Primary authors : Prof. RICHTER, Werner (iThemba LABS, Somerset West)

Co-authors :

Presenter : Prof. RICHTER, Werner (iThemba LABS, Somerset West)

Session classification : NPRP

Track classification : Track B - Nuclear, Particle and Radiation Physics

Type : Oral Presentation