



Contribution ID: 184

Type: Oral Presentation

Forbidden gap regions in ion-acoustic solitons

Wednesday, 1 July 2015 11:10 (20 minutes)

Abstract content (Max 300 words)
Formatting & Special chars

Plasma models composed of one and/or two (different mass) species of inertial ions and one and/or two (different temperature) species of (inertialess) electrons (isothermal and/or non-thermal) will be considered to investigate

forbidden gap regions (stopbands) in velocity space where ion-acoustic solitons do not propagate. It has been previously found [1] that these forbidden gap regions in velocity space occur between two passband regions which support the propagation of ion-acoustic solitons. The focus of the study will be to establish which plasma models favour the existence of stopband regions but also to determine how the sizes of the stopband regions are a function of the plasma parameters.

[1] Stopbands in the existence domains of acoustic solitons, F. Nsengiyumva, M. A. Hellberg, F. Verheest and R. L. Mace. Phys. Plasmas 21, 102301 (2014).

Apply to be considered for a student award (Yes / No)?

N/A

Level for award (Hons, MSc, PhD, N/A)?

N/A

Main supervisor (name and email) and his / her institution

N/A

Would you like to submit a short paper for the Conference Proceedings (Yes / No)?

No

Please indicate whether this abstract may be published online (Yes / No)

Yes

Primary author: Dr MAHARAJ, Shimul Kumar (South African National Space Agency (SANSA), Space Science Directorate)

Co-author: Prof. BHARUTHRAM, Ramesh (University of the Western Cape)

Presenter: Dr MAHARAJ, Shimul Kumar (South African National Space Agency (SANSA), Space Science Directorate)

Session Classification: Space

Track Classification: Track D2 - Space Science