



Contribution ID: 197

Type: Oral Presentation

## Decomposition of the field within an apertured plano-concave resonator

*Wednesday, 13 July 2011 11:15 (15 minutes)*

A method of experimentally decomposing a field within an apertured plano-concave resonator cavity into a forward propagating field (plane to concave) and a backward propagating field (concave to plane) is presented. The use of an intra-cavity aperture which is a simple method of laser beam shaping is a means of mode selection as higher-order modes are actively discriminated. This apertured cavity is modelled by two fundamental resonator theories, namely, Fox-Li analysis and Laguerre-Gaussian decomposition where they are used in the determination of the respective beam profiles at a specific plane. These theories traditionally differ from one another for an apertured cavity where the Gaussian beam within the cavity is truncated. A preliminary set-up is characterized for Gaussian propagation in an attempt to verify that the cavity is viable and accurate. A comparison of experimental data with the theories is presented.

**Level (Hons, MSc, <br> &nbsp; PhD, other)?**

MSc

**Consider for a student <br> &nbsp; award (Yes / No)?**

Yes

**Would you like to <br> submit a short paper <br> for the Conference <br> Proceedings (Yes / No)?**

No

**Primary author:** Mr NAIDOO, Darryl (Council for Scientific and Industrial Research)

**Co-authors:** Prof. FORBES, Andrew (Council for Scientific and Industrial Research); Prof. AÏT-AMEUR, Kamel (CIMAP-ENSICAEN)

**Presenter:** Mr NAIDOO, Darryl (Council for Scientific and Industrial Research)

**Session Classification:** LOS

**Track Classification:** Track C - Lasers, Optics and Spectroscopy