



Contribution ID: 71

Type: Oral Presentation

On formation of bubble structure near the island of inversion

Tuesday, 12 July 2016 11:40 (20 minutes)

Abstract content
 (Max 300 words)
Formatting &
Special chars

Depletion of the central density or formation of a hollow structure (Bubble) drew a lot of interest both from the theorists [1-3] as well as from the experimentalists [4,5] due to its conflicting nature vis-a-vis the property of saturation of the nuclear force. One can expect gross deviation from density saturation for finite Fermi systems in which single particle spectrum is discrete instead of a continuum. The formation of shells changes drastically for exotic nuclei having large isospins. As a result new properties of nuclei manifest in this region.

It is generally believed that non-occupation of s-states of exotic nuclei leads to central depression of density.

References:

- [1] X. Campi and D. W. L. Sprung, Phys. Lett. B46, 291 (1973).
- [2] M. Bender et al, Phys. Rev. C 60, 034304 (1999).
- [3] B. G. Todd-Rutel et al J, Phys. Rev. C69, 021301(R) (2004).
- [4] E. Khan et al, Nucl. Phys. A800, 37 (2008).
- [5] L. Gaudefroy et al., Phys. Rev. Lett. 97, 092501 (2006).
- [6] R. Bhattacharya, Nucl. Phys. A913, 1 (2013)
- [7] D.T. Yordanov et al, Phys. Rev, Lett. 108,042504 (2012)

Primary author: Prof. BHATTACHARYA, Rupayan (University of Calcutta)

Presenter: Prof. BHATTACHARYA, Rupayan (University of Calcutta)

Session Classification: Parallel Track B

Track Classification: High Energy, Nuclear and Particle physics