



Contribution ID: 292

Type: Oral Presentation

Development of a large mode area (LMA), high-power, thulium-doped fibre laser

Tuesday, 5 July 2016 10:00 (20 minutes)

Abstract content (Max 300 words) **Formatting & Special chars**

Thulium-doped fibre lasers are useful 2 micron sources that can be used for light detection and ranging (LIDAR), free space propagation of light, medical surgery and pumping of optical parametric oscillators (OPO). We investigated a high-power, large mode area (LMA), free space coupled, thulium-doped fibre laser which was pumped by 793 nm laser diodes. The fibre laser was characterized with regard to efficiency, beam quality, output power stability, and wavelength. Characterisation was also done with two intra-cavity volume Bragg gratings (VBG's). We achieved more than 60 W of stable, continuous wave output from a 3.7 m long thulium-doped fibre with a near theoretical limit slope efficiency of 54%. This output was achieved at the free running output wavelength of 2014 nm. The beam profile had a measured beam quality factor (M2) of 1.85 at 60 W of average laser power.

Apply to be considered for a student award (Yes / No)?

Yes

Level for award (Hons, MSc, PhD, N/A)?

MSc

Main supervisor (name and email) and his / her institution

Dr. Lorinda Wu, LWua@csir.co.za, Council for Scientific and Industrial Research (CSIR)

Would you like to submit a short paper for the Conference Proceedings (Yes / No)?

No

Please indicate whether this abstract may be published online (Yes / No)

Yes

Primary author: Mr MORRIS, Daniel (Council for Scientific and Industrial Research)

Co-authors: Prof. FORBES, Andrew (CSIR); Dr WU, Lorinda (CSIR-NLC); Dr KOEN, Wayne (CSIR National Laser Centre)

Presenter: Mr MORRIS, Daniel (Council for Scientific and Industrial Research)

Session Classification: Photonics

Track Classification: Track C - Photonics