



Contribution ID: 231

Type: **Poster Presentation**

Dust heating by Alfvén waves using non-Maxwellian distribution function

Wednesday, 6 July 2016 16:10 (1h 50m)

Abstract content (Max 300 words) Formatting & Special chars

Quasilinear theory is employed in order to evaluate the resonant heating rate by Alfvén waves, of multiple species dust particles in a hot, collisionless, and magnetized plasma, with the underlying assumption that the dust velocity distribution function can be modeled by a generalized (r, q) distribution function. The kinetic linear dispersion relation for the electromagnetic dust cyclotron Alfvén waves is derived, and the dependence of the heating rate on the magnetic field, mass, and density of the dust species is subsequently investigated. The heating rate and its dependence on the spectral indices r and q of the distribution function are also investigated. It is found that the heating is sensitive to negative value of spectral index r .

Apply to be considered for a student award (Yes / No)?

No

Level for award (Hons, MSc, PhD, N/A)?

PhD

Would you like to submit a short paper for the Conference Proceedings (Yes / No)?

Yes

Please indicate whether this abstract may be published online (Yes / No)

Yes

Primary author: Dr KIRAN, Zubia (GC University, Lahore, 54000, Pakistan)

Presenter: Dr KIRAN, Zubia (GC University, Lahore, 54000, Pakistan)

Session Classification: Poster Session (2)

Track Classification: Track G - Theoretical and Computational Physics