

A New Horizon of Few-Body Problems: Exact Coulomb treatment and the energy-momentum translation of the three-body Faddeev equation

S. Oryu^a, T. Watanabe^a, Y. Hiratsuka^a, A. Kodama^a,
Y. Togawa^b, and M. Takeda^b

a) Dept. of Physics, Faculty of Science and Technology,
Tokyo University of Science, Noda Chiba 278-8510, Japan

b) Dept. of Information Sciences, Faculty of Science and Technology,
Tokyo University of Science, Noda Chiba 278-8510, Japan

E-mail: oryu@rs.noda.tus.ac.jp

Abstract. We propose an off-shell Coulomb-like amplitude, with an on-shell phase shift that is accurate to within 9-10 digits. The full Coulomb amplitude is separated into the leading and the auxiliary amplitudes using “the two-potential theory”. The leading amplitude reproduces most of the on- and off-shell parts, while the auxiliary amplitude contributes mainly to the off-shell part and minimally to the on-shell part.

Next we review in a critical manner, the three-body calculation method, developed during the last four decades. A reminder to a threshold behavior investigation method is pointed out, based on the three-body Faddeev equation. We discuss the Efimov physics and some extensions, which are recovered from our predictions. The method may suggest a promising technique to resolve the existing discrepancies between the current experimental and theoretical values.

1. Introduction

The Coulomb phase shift renormalization method has a long history [1],[2]-[7]. The most plausible method was presented by Alt et al. [8]-[11]. Recently, we checked the reliability of the renormalization method. It was found that the phase shift can be definitely represented by the renormalization method. However, the screening range is unusually long [12]. In order to obtain the proton-proton phase shift $\sigma_0(k)=96.8$ deg. at $E=1$ keV by the renormalization method, we need to solve the Lippmann-Schwinger (LS) equation in momentum space with the screening range of 3nm, 1400 Gauss points, to within 250 significant digits (see Figure 1). Unfortunately, a serious cancelation of significant digits occurs without any warning signs, misleading so the converges to a wrong, unreliable value. Therefore, we confirmed that the phase shift renormalization is well performed by such a proper significant digits. However, Alt’s method cannot treat the nuclear interaction by the usual FORTRAN program. We have presented the following generalized screened Coulomb potential $V^R(r)$ and the renormalization phase $\phi(k, R)$ at the Groningen conference [13]:

$$V^R(r) = \frac{2k\eta(k)}{r} e^{-(r/R)^m} \quad (1)$$

$$\phi(k, R) = \eta(k) \left(\ln 2kR - \gamma/m \right). \quad (2)$$

In this case, we obtain a short cut-off potential by increasing m . After that, Deltuva et al. used an effective short-range, screened Coulomb potential, by adopting $m = 5$. In the three-body calculation, such two-body information is required for a wide two-body “sub-energy” region $-\infty < z'' \leq E$. This is because,

$$z'' = E - \frac{q''^2}{2\mu} \quad (3)$$

$$\text{for } 0 \leq q'' < \infty. \quad (4)$$

In order to obtain the two-body off-shell amplitude for the screened Coulomb potential, by the LS equation, one needs zero energy. Therefore, the most remarkable region of the Coulomb off-shell amplitude is the origin i.e. $z = 0$ for the two- and three-body problems. The appropriate ranges in the energy region from 0.1keV to 100 MeV, are listed in Table 1.

Table 1. Screening ranges for $m = 1$ and $m = 5$ with respect to the generalized screened Coulomb potential by Eq.(1). These ranges are obtained so to guarantee the three digit accuracy in the phase shift. Present ranges belongs to the asymptotic region. It is found that the $m = 5$ case yields on the average one order of magnitude shorter range than the case $m = 1$. In order to obtain higher accuracy in the phase shift, still longer ranges are required. If the renormalization method is used, efforts for getting the shorter ranges were shown to be ineffective.

Energy [MeV]	$m = 1$	$m = 5$	σ_0 [deg.]
100	11 fm	4 fm	-0.370
50	35 fm	8 fm	-0.523
20	110 fm	17 fm	-0.826
10	300 fm	35 fm	-1.17
1	900 fm	620 fm	-3.66
0.1	1.0×10^4 fm	3.7×10^3 fm	-10.7
0.01	1.0×10^5 fm	7.7×10^4 fm	-16.3
0.001	3.0×10^6 fm	4.5×10^5 fm	96.8
0.0001	3.0×10^7 fm	6.2×10^6 fm	949.9

Despite of the suggestion of Deltuva et al., it is found that the case $m = 5$ does not essentially improve over $m = 1$ case [14]-[16]. Therefore, we are afraid that the general use of unsatisfied Coulomb off-shell amplitude could mislead the three-nucleon scattering calculation. So, one may come to despair considering that the ASZ method may not have been successful for the past four decades.

2. On-shell Coulomb Amplitude

We introduced an off-shell Coulomb-like amplitude in the last EFB-23 conference, in Aarhus Denmark 2016 [12]. Before that we have introduced so-called “critical range” which satisfies zero renormalization phase [17]-[25]. In this paper, we will review our method for the on-shell Coulomb amplitude where the Coulomb phase shift is defined by the screened Coulomb phase




Figure 1. Necessary digits to obtain three digits accurate Coulomb phase shift by the renormalization method with the LS equation for $m = 1$ case.




Figure 2. Necessary screening range to represent the three digits accurate Coulomb phase shift by the renormalization method. $m = 1$ case in Eq.(1).

shift plus a renormalization phase: $\sigma_0(k) = \delta_0^R(k) + \phi(k, R)$. Therefore, the on-shell scattering amplitude is separated into two parts for the S-wave case by

$$f_0^C(k) = e^{2i\phi(k,R)} \left(\frac{e^{2i\delta_0^R(k)} - 1}{2ik} \right) + \left(\frac{e^{2i\phi(k,R)} - 1}{2ik} \right). \quad (5)$$

However, the ‘‘renormalized Coulomb amplitude’’ is usually defined by the first term of the right hand side of Eq.(5), where the second term has been neglected without explanation [9].

In our method, we require $\phi(k, R) = 2n\pi$ for $n = 0, 1, 2, \dots$. Using this criterion, a ‘‘critical range’’ is obtained, and the screened Coulomb phase shift becomes the Coulomb phase shift itself: $\delta_0^R(k) = \sigma_0(k)$, and the on-shell amplitude is given by $f_0^C(k) = (e^{2i\delta_0^R(k)} - 1)/2ik$. In other words, the Coulomb phase shift can be obtained directly from the critical range. In FB20 conference (Chicago 2015), we pointed out that the screening range is not obtained by increasing




Figure 3. Necessary screening range to represent the three digits accurate Coulomb phase shift by the renormalization method. $m = 5$ case in Eq.(1).

a single range but by concatenating many discrete bands [26]. Since Alt's renormalization phase is given by Eq.(2) in the asymptotic region [27], we may set:

$$\phi(k, R) = \eta(k) \left(\ln 2kR - \frac{\gamma}{m} \right) = 2n\pi. \quad (6)$$

Therefore, our “critical range”, hereafter Alt's range, becomes:

$$R = \frac{1}{2k} \exp \left(\frac{2\pi}{\eta(k)} + \frac{\gamma}{m} \right). \quad (7)$$

We conclude that the screened Coulomb phase shift becomes the Coulomb phase shift by using this critical range. However, a very long range is required to satisfy the asymptotic properties (see Figures 2,3). Such a long range, due to the cancellation of significant digits, renders the solution of the LS equation quite hard.

In order to avoid these difficulties, we obtained the best range parameters, by a fitting method at a region much shorter than the asymptotic domain where the phase shift calculation guarantees nine to ten digits accuracy (see Figures 1-3, and Table 1) [27].

Hereafter, the term “critical range” will be used to refer to these best range parameters.

3. A new Method for the off-shell Coulomb-like Amplitude

We presented an off-shell Coulomb-like amplitude in the EFB23 conference [12]. In this paper, we review it and extend the results. We would like to introduce the two-potential theory in r -space. The Schrödinger equation for the Coulomb wave function may be written in operator form as:

$$(E - H_0 - V^{\mathcal{R}})|\psi^C\rangle = V^\phi|\psi^C\rangle = (V^C - V^{\mathcal{R}})|\psi^C\rangle. \quad (8)$$

Now, we assume a relation: $V^C|\psi^C\rangle \approx V^{\mathcal{R}'}|\psi^{\mathcal{R}'}\rangle$, where $V^{\mathcal{R}'}$ is a screened Coulomb potential of range \mathcal{R}' with proper asymptotic properties and $|\psi^{\mathcal{R}'}\rangle$ the corresponding wave function. This assumption makes sense in the frame of the above discussion of Alt's Coulomb renormalization method. Therefore, the asymptotic behavior for such a long range region leads

to $V^\phi|\psi^C \rangle \approx (V^{\mathcal{R}'} - V^{\mathcal{R}})|\psi^{\mathcal{R}'} \rangle \equiv V^{\phi'}|\psi^{\mathcal{R}'} \rangle \rightarrow 0$ (for $r \rightarrow \infty$). With this condition, we can define a general solution $|\psi^{\mathcal{R}} \rangle$ for the homogeneous equation in Eq.(8). Finally, the general solution of Eq.(8) is given by

$$|\psi^C \rangle = |\psi^{\mathcal{R}} \rangle + G^{\mathcal{R}}V^\phi|\psi^C \rangle \quad (9)$$

$$G^{\mathcal{R}} = [E - H_0 - V^{\mathcal{R}}]^{-1} = G_0 + G_0T^{\mathcal{R}}G_0 \equiv G_0\bar{\omega}^{\mathcal{R}}, \quad (10)$$

$$\bar{\omega}^{\mathcal{R}} = 1 + T^{\mathcal{R}}G_0, \quad (11)$$

$$\omega^{\mathcal{R}} = 1 + G_0T^{\mathcal{R}}, \quad (12)$$

where $G^{\mathcal{R}}$ is Green's function with the screened Coulomb potential $V^{\mathcal{R}}$.

By multiplying Eq. (9) from the left with V^ϕ , an amplitude $t^{\phi\mathcal{R}}$ may be defined using $|\psi^{\mathcal{R}} \rangle$ and the plane wave $|\psi_0 \rangle$, as:

$$V^\phi|\psi^C \rangle \equiv t^{\phi\mathcal{R}}|\psi^{\mathcal{R}} \rangle = t^{\phi\mathcal{R}}\omega^{\mathcal{R}}|\psi_0 \rangle \quad (13)$$

So, $t^{\phi\mathcal{R}}$ introduces all the long range properties in the Coulomb wave function, and also merits for the proper asymptotic behavior mentioned above.

Therefore, Eq.(9) is rewritten by using Eq.(10), and Eq.(13),

$$|\psi^C \rangle = |\psi^{\mathcal{R}} \rangle + G^{\mathcal{R}}t^{\phi\mathcal{R}}|\psi^{\mathcal{R}} \rangle = |\psi^{\mathcal{R}} \rangle + G_0\bar{\omega}^{\mathcal{R}}t^{\phi\mathcal{R}}\omega^{\mathcal{R}}|\psi_0 \rangle \quad (14)$$

$$= |\psi^{\mathcal{R}} \rangle + G^{\mathcal{R}}V^\phi|\psi^C \rangle = |\psi^{\mathcal{R}} \rangle + G_0\bar{\omega}^{\mathcal{R}}V^\phi|\psi^C \rangle. \quad (15)$$

Multiplying Eq.(14) from the left with V^ϕ , and using Eq.(13), yields:

$$V^\phi|\psi^C \rangle = t^{\phi\mathcal{R}}|\psi^{\mathcal{R}} \rangle = V^\phi|\psi^{\mathcal{R}} \rangle + V^\phi G^{\mathcal{R}}t^{\phi\mathcal{R}}|\psi^{\mathcal{R}} \rangle. \quad (16)$$

Here, we define a Coulomb-like off-shell T-matrix \tilde{T}^C by

$$G_0^{-1}|\psi^C \rangle = V^C|\psi^C \rangle \equiv \tilde{T}^C|\psi_0 \rangle, \quad (17)$$

$$G_0^{-1}|\psi^{\mathcal{R}} \rangle = V^{\mathcal{R}}|\psi^{\mathcal{R}} \rangle = T^{\mathcal{R}}|\psi_0 \rangle, \quad (18)$$

where \tilde{T}^C has all the properties of the long range Coulomb wave function and the potential. In addition it satisfies the Coulomb asymptotic behavior and our requirement of Eq.(6) which is equivalent to the *Lemma* stated in our recent article [26].

Multiplying Eq.(14) from the left with V^C and using Eq.(17), Eq.(18), we obtain

$$\tilde{T}^C|\psi_0 \rangle = T^{\mathcal{R}}|\psi_0 \rangle + \bar{\omega}^{\mathcal{R}}t^{\phi\mathcal{R}}\omega^{\mathcal{R}}|\psi_0 \rangle \quad (19)$$

$$= T^{\mathcal{R}}|\psi_0 \rangle + \bar{\omega}^{\mathcal{R}}V^\phi|\psi^C \rangle. \quad (20)$$

Therefore, Eq. (19), Eq. (16), and Eq. (10) lead to the following operator relations:

$$\tilde{T}^C = T^{\mathcal{R}} + T^{\phi\mathcal{R}} = T^{\mathcal{R}} + \bar{\omega}^{\mathcal{R}}t^{\phi\mathcal{R}}\omega^{\mathcal{R}}, \quad (21)$$

$$t^{\phi\mathcal{R}} = V^\phi + V^\phi G^{\mathcal{R}}t^{\phi\mathcal{R}} \quad (22)$$

$$T^{\mathcal{R}} = V^{\mathcal{R}} + V^{\mathcal{R}}G_0T^{\mathcal{R}}, \quad (23)$$

where Eq. (21) is known as the amplitude by the two-potential theory. Multiplying from the left Eq.(20) with $\langle \psi_0 |$ and using $\langle \psi_0 | \bar{\omega}^{\mathcal{R}} = \langle \psi^{\mathcal{R}} |$, we obtain

$$\begin{aligned} \langle \psi_0 | \tilde{T}^C | \psi_0 \rangle &= \langle \psi_0 | T^{\mathcal{R}} | \psi_0 \rangle + \langle \psi_0 | \bar{\omega}^{\mathcal{R}} V^\phi | \psi^C \rangle \\ &= \langle \psi_0 | T^{\mathcal{R}} | \psi_0 \rangle + \langle \psi^{\mathcal{R}} | V^\phi | \psi^C \rangle. \end{aligned} \quad (24)$$

Eq. (24) is equivalent to Eq. (21), and the amplitude $T^{\phi\mathcal{R}}$ is obtained by using the partial wave functions $|\psi_L^{\mathcal{R}}\rangle$ and $|\psi_L^C\rangle$.

$$T_L^{\phi\mathcal{R}}(p, p'; E) \equiv \langle \psi_L^{\mathcal{R}} | V^\phi | \psi_L^C \rangle = \int_0^\infty \psi_L^{\mathcal{R}*}(p, r) V^\phi(r, k) \psi_L^C(p', r) r^2 dr. \quad (25)$$

This result guarantees the existence of the off-shell Coulomb-like T-matrix by Eq. (21) and Eq. (25), which was not clearly stated in [28]. The off-shell value given by Eq.(25), is calculated numerically, where the integrand is highly oscillatory and vanishes for very large values of r . Therefore, Eq.(25) contributes to off-shell, and half-on-shell values. In our method, the on-shell value of $T^{\mathcal{R}}$ in Eq.(21) gives 9–10 digits of accuracy [27]. Therefore, it is obvious that the on-shell value of $T^{\phi\mathcal{R}}$ contains only the next order correction need to obtain an accuracy of 9–10 digits in the phase shift.

However, the numerical calculation of the on-shell value of Eq.(25) is very sensitive to the convergence condition. Therefore, in order to calculate the on-shell value of Eq.(25), we can adopt a screened Coulomb potential with a very large range ($\mathcal{R} \ll \mathcal{R}'$) which appears in the asymptotic region and is introduced to obtain the solution of Eq.(8). For such an \mathcal{R}' , we can accomplish the following relation:

$$T_L^{\phi\mathcal{R}}(k, k; E) = \int_0^\infty \psi_L^{\mathcal{R}*}(k, r) V^\phi(r, k) \psi_L^C(k, r) r^2 dr \equiv \Delta_L \quad (26)$$

$$\rightarrow \int_0^\infty \psi_L^{\mathcal{R}*}(k, r) V^{\phi'}(r, k) \psi_L^{\mathcal{R}'}(k, r) r^2 dr \cong -\frac{|T_L^{\mathcal{R}}|^2}{\mathcal{I}m(T_L^{\mathcal{R}})} \times \phi'_L(k) \quad (27)$$

with

$$\phi'_L(k) \equiv \sigma_L(k) - \delta_L^{\mathcal{R}}(k) = \tan^{-1} \frac{\mathcal{I}m(T_L^{\mathcal{R}} + \Delta_L)}{\mathcal{R}e(T_L^{\mathcal{R}} + \Delta_L)} - \tan^{-1} \frac{\mathcal{I}m(T_L^{\mathcal{R}})}{\mathcal{R}e(T_L^{\mathcal{R}})}, \quad (28)$$

where the last term in Eq.(27) is an auxiliary amplitude, $\phi'_L(k)$ is the discrepancy between the analytic Coulomb phase shift and our direct phase shift. We can find an adequate \mathcal{R}' which satisfies Eq.(27). In ref.[12], we mentioned that the value of Eq.(26) vanishes. However, such a condition is satisfied only for the case where the eigen-values of $V^C|\psi^C\rangle$ and $V^{\mathcal{R}}|\psi^{\mathcal{R}}\rangle$ are identical. Our fitting procedure albeit not exact, reproduces the Coulomb phase shift with an accuracy of 9–10 digits.

Therefore, we can conclude that the first term $T^{\mathcal{R}}$ represents almost all on- and off-shell Coulomb amplitude, with a compensation coming from the off-shell part of the second term.

4. Review of the three-hadron problem

Half a century has passed since the Faddeev equation was presented [29]. In the beginning, it seemed that the obtained results fitted the experimental data well, in contrast to the results of other methods. After that, improved results have been mainly due to the rapid growth of the computer capacity as to how many partial waves can be incorporated, and what kind of realistic two-body and three-body forces can be treated. These efforts have been done somewhat with a technical professionalism. However, the ideas of the theoretical and methodological approach of the Faddeev equation were investigated only in the beginning, that is, in 1960s and 1970s [30]-[34]. It seems that today, is a good time to start a “deep critical rethinking” of the results obtained so far by the usual Faddeev calculations, in order to understand whether the fundamental problems of the Faddeev equation still exist or not.

First of all, the quantitative agreements for A_y are not yet satisfied [35]. Recently, the intensive discussion about the disagreements for the cross section minimum and A_y maximum

were pointed out by Sagara [36]. The so-called ‘‘Sagara discrepancy’’ was illustrated by a ratio between theoretical and experimental data in the cross section minimum and the maximum of A_y : $S_{\text{disc}} \equiv (\text{calculation-experiment})/\text{experiment}$ [37][38]. Such discrepancies have an energy dependence which is commonly inversely proportional to the energy. It was claimed that the cross section minimum problem was solved by using a three-body force named $2\pi 3\text{NF}$ [39]. However, it seems that the A_y puzzle has not yet been solved by such a systematic way with $2\pi 3\text{NF}$ and some other intensive 3NFs [40]. We are concerned about the energy dependence of the Sagara discrepancy not only for the A_y puzzle, but also the cross section minimum. If such a 3NF can fundamentally reproduce the cross section minimum, the A_y -puzzle should also be solved by the 3NF at the same time. However, the puzzle is not solved yet. The concern arises from whether the 3NF is fundamental or an ad hoc remedy in the three-nucleon problem.

Another problem discussed but remained unsolved is the break-up problem in the case of ‘‘the Space star anomaly’’, although 3NF and many different so-called realistic potentials were used. Also an incomplete differential cross section could not yet be represented by the Faddeev calculation [30].

On the other hand, the discrepancy in the triton binding energy between the experimental data is compensated by a three-body force [41]. It was found that the doublet n-d scattering length (a_{nd}^2) is closely related to the three-body force[42]. The closed relation between a_{nd}^2 and 3NF is reminiscent of the relation between the ‘‘n-d threshold’’ and the 3NF. The n-d threshold is not clearly defined in the Faddeev equation unfortunately, because the Born term of AGS equation or the ‘‘n-d effective potential’’ does not diverge at the n-d threshold.

Many attempts are carried out by using realistic potentials and corresponding three-body forces, relativistic effects, the Coulomb interaction, new degrees of freedom etc. However, those methods could not satisfactorily explain the above discrepancies.

The three-body force is mainly an isotropic interaction. Furthermore, the renormalization phase of the Coulomb amplitude is also isotropic. Such isotropic interactions can not affect the P-wave dominant A_y value, and the effect of the isotropic interaction could work somewhat negatively. Therefore, we suspect that the energy dependence in the Sagara discrepancy could be closely related to an ‘‘energy-momentum usage or kinematics’’ in the Faddeev equation.

The second is the three-body bound state which is obtained by solving the two-variable eigen-equation as the three-body Faddeev equation. However, the two-variable eigen-equation is ‘‘not well defined’’ mathematically, although the one variable eigen-equation is well defined [43].

Therefore, we should separate the two-variable three-body Faddeev equation into one-variable two-body LS equations, and a one-variable quasi-two-body multi-channel LS equation. Each of them gives eigen-values $\epsilon^{(2)}$ and ζ by solving one-variable eigen-equations, where we obtain a separation energy ζ for the reaction $(abc) \rightarrow (bc) + a$ by the quasi-two-body multi-channel LS equation. Therefore, the three-body binding energy is obtained by $\epsilon^{(3)} = \zeta + \epsilon^{(2)}$ [44].

This method coincides with the Jacobi-coordinate anti-symmetrized molecular dynamics, where bound colonies are grown up sequentially from many free-particle groups, by cooling the system in a timely manner, and not instantaneously as in the usual Faddeev method. [45]-[49].

4.1. Three-body kinematics

Now, let us start with the well-known three-body kinematics for the scattering with a incident particle and a target. The incident particle has a kinetic energy E_{Lab} with the momentum q_L ,

$$E_{\text{Lab}} = \frac{q_L^2}{2m} = \frac{q_L^2}{2(m + M_t)} + E_{\text{cm}} = \left(\frac{m}{m + M_t}\right)E_{\text{Lab}} + E_{\text{cm}} \quad (29)$$

$$E = E_{\text{cm}} - \epsilon_B = \frac{q^2}{2\mu} + \frac{p^2}{2\nu}, \quad (\text{target binding energy} = \epsilon_B) \quad (30)$$

$$E_{\text{cm}} = \frac{\bar{q}^2}{2\mu} + \frac{\bar{p}^2}{2\nu}, \quad (\text{target unbound system}), \quad (31)$$

where m and M_t are the masses of the incident particle and the target. μ denotes the reduced mass between a pair and the spectator particle, and ν is the reduced mass in the pair, respectively. Therefore, q is the relative momentum between the pair and the spectator, and p is the relative momentum in the pair after three-body break-up. E_{cm} is the three-body center of mass energy, and ϵ_B is the target binding energy. Therefore E is the three-body free energy after the deuteron break-up. If there are no two-body sub-bound states in the intermediate state, then the three-body total energy E_{cm} should be used, where one of the three particles, at least, is virtual.

Usually, the Faddeev equation and the AGS equation commonly use the three-body free energy E in the Green's function: $G_0(E)$ where real-three-particles are transferred in between two different three-body channels. However, if both of the three-body channels are deuteron bound states, the spectator particles gain the deuteron binding energy. Therefore, the initial and the final state's energies are three-body total energies E_{cm} . In the Born term of AGS equation, the Green's function should be written by $\bar{G}_0(E_{\text{cm}})$, because all the particle-lines in the Feynman diagram are "outer-lines" where the energy-momentum should be conserved. Therefore, a transferred momentum in the Green's function should be partly virtual.

Furthermore, $E_{\text{cm}} = 0$ indicates the quasi-two-body threshold in the three-body scattering which causes a divergence of the Born term. This means that the n-d scattering length is calculated with such analytic properties as to $E_{\text{cm}} = 0$, although the original Faddeev equation has no singularity at this threshold: $E = -\epsilon_B$. We can call such a critical energy point at $E_{\text{cm}} = 0$ the "quasi-two-body threshold" (Q2T) in the three-body system.

On the other hand, in the intermediate three-body lines as the "inner lines" of the Feynman diagram, we are concerned about two-type of three-body energies: the three-body free energy E and the three-body total-energy E_{cm} . Therefore, the Green's function which is sandwiched between the bound state and unbound states could be distinguished by $G_0(E)$ and $\bar{G}_0(E_{\text{cm}})$.

Therefore, the Born term of the quasi-two-body equation for N - d system could be written by \bar{q} and \bar{q}' which are represented by the relative momentum between a pair and the spectator.

We can easily introduce a new Born term by using "an energy-momentum translation" in the usual Born term,

$$E \rightarrow E_{\text{cm}} = E + \epsilon_B \quad (32)$$

$$q^2/2\mu \rightarrow \bar{q}^2/2\mu = q^2/2\mu + \epsilon_B, \quad (33)$$

$$\bar{p}^2/2\nu = p^2/2\nu, \quad (34)$$

we obtain

$$Z_{\alpha,\beta}(q, q'; E) = \frac{g_\alpha(p)g_\beta(p')(1 - \delta_{\alpha\beta})}{(E + \epsilon_B) - (q^2/2\nu_\beta - q'^2/2\nu_\alpha - qq'x/m + \epsilon_B)} \quad (35)$$

$$= \frac{g_\alpha(\bar{p})g_\beta(\bar{p}')(1 - \delta_{\alpha\beta})}{E_{\text{cm}} - \bar{q}^2/2\nu_\beta - \bar{q}'^2/2\nu_\alpha - \bar{q}\bar{q}'\bar{x}/m} = Z_{\alpha,\beta}(\bar{q}, \bar{q}'; E_{\text{cm}}). \quad (36)$$

where $x = \mathbf{q}\mathbf{q}'/qq'$ and $\bar{x} = \bar{\mathbf{q}}\bar{\mathbf{q}}'/\bar{q}\bar{q}'$. The propagator becomes,

$$\tau_\gamma(E - q'^2/2\mu) = \tau_\gamma(E_{\text{cm}} - \bar{q}'^2/2\mu). \quad (37)$$

Therefore, the energy-momentum translation does not affect the function itself, however the integral may be modified due to the prolonged range.

4.2. Non-Efimov but Efimov-like states

One author (S.O.) pointed out that $E_{\text{cm}} = 0$ in Eq.(36) for NN π system, generates a singularity-coincidence between the Born term and the propagator $\tau_\gamma(E_{\text{cm}} - \bar{q}^2/2\mu)$ [50]. This coincidence is very similar to the Coulomb scattering in the LS equation where the singularity of the Coulomb potential and of the propagator occurs simultaneously.

Since our potential singularity in the present case arises at $E_{\text{cm}} = 0$, then we carried out an energy average around $E_{\text{cm}} = 0$ after the Fourier transformation of Eq.(36). Finally, we obtained $V(r) = V_0 a^{2\gamma+2}/[r(r/2 + a)^{2\gamma+2}]$ -type potentials with parameters a and γ and a depth constant $V_0 (< 0)$ for the lowest order term in the local potential expansion of the Born term. Therefore, we obtained $V_0(2a)/r^2$ potential for $\gamma = -1/2$, and Van der Waals potentials $V_0(2a)^5/r^6$ for $\gamma = 3/2$, $V_0(2a)^6/r^7$ for $\gamma = 2$ potentials in the long range region for $a \ll r$. While in the shorter range region $r \ll a$, they become Yukawa-type potentials $V_0 e^{-r/2a}/r$, $V_0 e^{-5r/2a}/r$, and $V_0 e^{-6r/2a}/r$, respectively (see Table 2). Beside the lowest order, the higher order terms affect the shorter range region adding to the Yukawa-type potential. Therefore, the shorter range region could be strongly modified from the Yukawa-type potential near the core region, however our longer range predicted potential can not be changed by the higher order terms.

These potentials are closely related to the mass ratio between a transfer particle and the parent particles, which can be easily understood by the reason that if the massless particle is transferred, then the potential will be $1/r$ which is given by $\gamma = -1$. It is interesting that $\gamma = -1$ or a massless particle creates a unique potential V_0/r where two-type of potentials for $a \ll r$ and $r \ll a$ cross each other (see Table 2).

The above discussion, that two types of potentials, a long range and a short range, arise from a particle-transfer mechanism, seems to be a universal structure in physics. If there is a counterargument about our “energy average” method, we can directly solve the three-body equation with Eq.(36), where we need a special technique to treat it in the neighborhood of the singular region. Finally, we can also obtain bound states near the E2Q-threshold or the Q2T.

It is well known that $1/r^2$ potential generates the Efimov-like states, although our case does not require the infinite number of the two-body scattering length: $a_{\text{scatt}} \neq \pm\infty$. However, when the two-body binding energy becomes zero then $0 = E_{\text{cm}} = E + \epsilon_B \rightarrow E$ which is the case of Efimov states [51][52]. In this sense, our prediction covers the Efimov states. We can say that our prediction at the Q2T is generalized to other thresholds such as three-body, four-body, five-body, ..., thresholds [53]. This fact suggests that “observed and unobserved energy levels” in any many-body system give rise to a long range potential near the threshold. We would like to suggest that these levels could be a key which can “control the nuclear reactions” where the long range attractive nuclear interaction could interfere with the Coulomb repulsive force in the pico-size region [50].

5. Conclusion and Discussion

The reliability of the Coulomb renormalization method is confirmed. However, its application to the three-body problem is almost impossible, because the screening range is too long. When Alt’s-type Coulomb renormalization is used, a serious cancelation of significant digits occurs in the two-body LS and in the three-body Faddeev equations. This cancelation causes the method to converge, without any warning, to an erroneous value, hence rendering the calculation unreliable. We present a shorter ranged screened Coulomb potential which can reproduce the Coulomb phase shift to within 9–10 digits accuracy. Furthermore, the off-shell Coulomb-like amplitude is proposed by the two-potential method in r -space representation where the off-shell amplitude can be calculated numerically. By our theory, the Coulomb treatment in the three-body problem could be enormously improved for the first time in a period of half a century.




Figure 4. Bound state energy levels for the potentials $1/r$ (left) and $1/r^2$ (right) where the ground state energies are fitted. The vertical unit is arbitrary but the logarithmic unit of the absolute value is taken.

Table 2. An effective potential $V_0 a^{2\gamma+2}/[r(r/2 + a)^{2\gamma+2}]$ is illustrated, which is given by an energy average around the Q2T (or $E_{\text{cm}} = 0$) with two-parameters a and γ . The potential properties for the longer and shorter ranges are shown with respect to the parameter γ . $V_0(<0)$, a potential depth which is analytically given by the Born term of the form Eq.(36).

γ	short range potential $r \ll a$	potential	long range potential $a \ll r$
-1	V_0/r	V_0/r	V_0/r
-1/2	$V_0 e^{-r/2a}/r$	$V_0(2a)/[r(r+2a)]$	$V_0(2a)/r^2$
0	$V_0 e^{-2r/2a}/r$	$V_0(2a)^2/[r(r+2a)^2]$	$V_0(2a)^2/r^3$
1/2	$V_0 e^{-3r/2a}/r$	$V_0(2a)^3/[r(r+2a)^3]$	$V_0(2a)^3/r^4$
1	$V_0 e^{-4r/2a}/r$	$V_0(2a)^4/[r(r+2a)^4]$	$V_0(2a)^4/r^5$
3/2	$V_0 e^{-5r/2a}/r$	$V_0(2a)^5/[r(r+2a)^5]$	$V_0(2a)^5/r^6$
2	$V_0 e^{-6r/2a}/r$	$V_0(2a)^6/[r(r+2a)^6]$	$V_0(2a)^6/r^7$
.
.

Finally, we would like to emphasize that the method is very general, and it can treat all kinds of charged particle systems, from the electron-electron to the heavy ion-heavy ion, using the universal variables.

We reviewed the discrepancy between the experimental data and the usual Faddeev calculated results. Sagara discrepancy is the most important admonition for this problem. We also added some other problems that seemed solvable by incorporating the three-body force. We pointed out that the Q2T is important for improving the Faddeev treatment. One of the examples was the case of the $NN\pi$ system, and we presented a possible existence of a local long-range NN' potential around Q2T, and NN' scattering length and πD one [54][53]. This potential results to some interesting bound states, and to the existence of long-range potentials such as the $1/r^2$ type and the Van der Waals potential. The $1/r^2$ potential gives infinite number of bound states below the Q2T, without Efimov's requirement for infinite scattering length of two-body sub-

system. Therefore, our theory predicts, in addition to the Efimov states, some other interesting aspects of nuclear physics.

Acknowledgments

The authors would like to pay tribute to the memory of the late Professor S. A. Sofianos, for his support and continuous encouragement since the very beginning of this idea.

References

- [1] Tobocman W and Kalos M H: 1955 Numerical Calculation of (d,p) Angular Distributions. *Phys. Rev.***97**, 132
- [2] Veselova A M: 1970 Separation of two-particle Coulomb singularities in a system of three charged particles. *Teor. Mat. Fiz.***3**, 326 (in Russian)
- [3] Veselova A M: 1970 Separation of two-particle Coulomb singularities in a system of three charged particles. *Theor. Math. Phys.***3**, 542 (Leningrad State University, Plenum Publishing Corporation)
- [4] Veselova A M: 1972 Determination of scattering amplitudes in problems with two and three charged particles. *Teor. Mat. Fiz.***13**, 368 (in Russian)
- [5] Veselova A M: 1972 Determination of scattering amplitudes in problems with two and three charged particles. *Theor. Math. Phys.***3**, 1200 Leningrad State University, Plenum Publishing Corporation
- [6] Veselova A M: 1978 Integral equations for three particles with Coulomb long-range interaction. *Teor. Mat. Fiz.***35**, 180 (in Russian)
- [7] Veselova A M: 1978 Integral equations for three particles with Coulomb long-range interaction (in Russian). *Theor. Math. Phys.***35**, 395 Leningrad State University, Plenum Publishing Corporation
- [8] Alt E O, Sandhas W, Zankel H and Ziegelmann H: 1976 Coulomb Corrections in Proton-Deuteron Scattering. *Phys. Rev. Lett.***37**, 1537
- [9] Alt E O, Sandhas W and Ziegelmann H: 1978 Coulomb effects in three-body reactions with two charged particles. *Phys. Rev.***C17**,1981
- [10] Alt E O and Sandhas W: 1980 Scattering amplitudes and integral equations for the collision of two charged composite particles. *Phys. Rev.***C21**, 1733
- [11] Alt E O, Sandhas W and Ziegelmann H: 1985. Calculation of proton-deuteron phase parameters including the coulomb force. *Nucl. Phys.***A445**, 429
- [12] Oryu S, Watanabe T, Hiratsuka Y and Togawa Y: 2017, A Coulomb-like Off-Shell T-matrix with the Correct Coulomb Phase Shift, *Few-Body Syst.* 58:95, DOI 10.1007/s00601-017-1258-6
- [13] Oryu S, Nishinohara S, Sonaoda K, Shiiki N, Togawa Y and Chiba S: 2005, Oryu, S: 2004, talk in EFB19 (Groningen): A novel three-charged-particle Faddeev type equation in momentum space, AIP Conference Proc. the 19-th European Conference on Few-Body Problems in Physics, Vol 768, 433-435.
- [14] Deltuva A, Fonseca A C and Sauer P U: 2005 Momentum-space description of three-nucleon breakup reactions including the Coulomb interaction *Phys. Rev.***C72**, 054004
- [15] Deltuva A, Fonseca A C, Kievsky A, Rosati S, Sauer P U and Viviani M: 2005 Benchmark calculation for proton-deuteron elastic scattering observables including the Coulomb interaction. *Phys. Rev.* **C71**, 064003
- [16] Deltuva, A, Fonseca A C and Sauer P U: 2005 Momentum-space treatment of the Coulomb interaction in three-nucleon reactions with two protons. *Phys. Rev.* **C71**, 054005
- [17] Hiratsuka Y, Oryu S and Gojuki S: 2013 D(p,p)D elastic scattering with rigorous Coulomb treatment. *J. Phys G: Nucl. Part. Phys.***40**, 025106 (17pp)
- [18] Oryu S, Hiratsuka Y, Nishinohara S and Chiba S: 2012 Proton-proton phase shifts calculations in momentum space by a rigorous Coulomb treatment. *J. Phys.G: Nucl. Part. Phys.***39**, 045101 (12pp)
- [19] Oryu S, Nishinohara S, Shiiki N, and Chiba S: 2007 Coulomb phase shift calculation in momentum space. *Phys. Rev.***C75**, 021001
- [20] Nishinohara S, Chiba S and Oryu S: 2007 The Coulomb scattering in momentum space for few-body systems. *Nucl. Phys.* **A790**, 277c
- [21] Oryu S: 2006 Two- and three-charged particle nuclear scattering in momentum space: A two-potential theory and a boundary condition model. *Phys. Rev.***C73**, 054001
- [22] Oryu S: 2007 Erratum: Two- and three-charged particle nuclear scattering in momentum space: A two-potential theory and a boundary condition model. *Phys. Rev.* **76** 069901
- [23] Nishinohara S: 2006 A study of Coulomb scattering by the momentum representation (in Japanese). Doctor thesis in Tokyo University of Science
- [24] Hiratsuka Y: 2014 A study of the rigorous solution for two- and three-charged nucleons scattering in momentum space (in Japanese). Doctor thesis in Tokyo University of Science

- [25] Oryu S, Hiratsuka Y and Watanabe T: 2016 A New Treatment Below the Three-Body Break up Threshold in the $NN\pi$ System *EPJ Web of Conferences* **113** 08014
- [26] Watanabe T, Hiratsuka Y and Oryu S: 2016 Effective Screened Coulomb Potential from e^-e^- to ^{208}Pb – ^{208}Pb Systems. *EPJ Web of Conf.* **113**, 03018
- [27] Watanabe T, Y. Hiratsuka, Oryu S, Togawa Y: 2017 A New Feature of the Screened Coulomb Potential in Momentum Space *Few-Body Syst.* 58:80; DOI 10.1007/s00601-017-1238-x,
- [28] H. van Haeringen: 1985 *Charged Particle Interactions Theory and Formulas* (Coulomb Press Leyden, Netherlands)
- [29] Faddeev L D: 1960 *Zh. Eksp. Theor. Fiz.* **39** 1459
Faddeev L D: 1961 *Sov. Phys. -JETP* **12** 1014
- [30] Aaron R and Amado R D: 1966 Theory of the Reaction $n+d \rightarrow n+n+p$. *Phys. Rev.* **150**, 857
- [31] Lovelace C: 1964 Practical Theory of Three-Particle States. I. Nonrelativistic. *Phys. Rev* **135**, B1225
- [32] Alt E O, Grassberger and Sandhas W: 1967 Reduction of the three-particle collision problem to multi-channel two-particle. Lippmann-Schwinger equations. *Nucl. Phys.* **B2**, 167
- [33] Cahill R T and Sloan I H: 1971 Theory of neutron-deuteron break-up at 14.4 MeV. *Nucl. Phys.*, **A165** 161
- [34] Ebenhöf W: 1972 The $n+d$ break-up reaction with separable potentials. *Nucl. Phys.* **191** 97
- [35] Koike Y and Heidenbauer J: 1987 The neutron-deuteron elastic scattering with the Paris potential. *Nucl. Phys.* **A463** 365c
- [36] Sagara K: 2010 Experimental Investigations of Discrepancies in Three-Nucleon Reactions *Few-Body Syst* 48:59-108 DOI 10.1007/s00601-010-0096-6
- [37] Takemiya N: 1985 On the low energy n - d scattering. *Prog. Theor. Phys.* 74, 301
Takemiya, N: 1991 Analyzing powers in elastic nucleon-deuteron scattering. *Prog. Theor. Phys.* 86, 975
- [38] Koike Y and Ishikawa S: 1998 Possible origin of Sagara discrepancy in the cross section minimum of p - d elastic scattering. *Nucl. Phys. A* 631, 683c
- [39] Witała H, et al.: 1998 Cross section minima in elastic Nd scattering: possible evidence for three-nucleon force effects. *Phys. Rev. Lett.* 81, 1183
- [40] Witała H, Golak J, Skibinski R, Topolnichki K, Kamada H, Epelbaum E, Glöckle W, Krebs H, Polyzou W N and Nogga A: 2013 Calculations of Three-Nucleon Reactions. *Few-Body Sys.* **54** 897
- [41] Ishikawa S, Sasakawa T, Sawada T, and Ueda T: 1984 Contribution of Three-Nucleon Potential to Triton Binding Energy. *Phys. Rev. Lett.* **53**, 1877
- [42] Oryu S and Yamada H: 1994 Effects of the three-body force in three-nucleon system *Phys. Rev* **C49** 2337
- [43] Furutani K, Private communication.
- [44] Oryu S, Hiratsuka Y, and Watanabe T: 2016 *EPJ Web of Conferences* 122, 08001 (2016) CNR*15, DOI: 10.1051/epjconf/201612208001, Few-Body Problem in Nuclear Reactions: *Beyond the horizon of the three-body Faddeev equations*
- [45] Watanabe T and Oryu S: 2006 A New Antisymmetrized Molecular Dynamics Approach to Few-Nucleon Systems. *Prog. Theor. Phys.* **116** 429-434
- [46] Watanabe T, Oosawa M, Saito K and Oryu S: 2009 A new molecular dynamics calculation and its application to the spectra of light and strange baryons. *J. Phys. G:* **36** 015001
- [47] Watanabe T, Oryu S and Saito K: 2009 NEW APPROACH FOR FEW-BODY SYSTEMS BASED ON JACOBI-COORDINATE BASIS ANTI-SYMMETRIZED MOLECULAR DYNAMICS FROM ATOMS TO QUARK-SYSTEMS. *Mod. Phys. Lett.* **A24** 797-803
- [48] Taniguchi Y, Watanabe T, Sawado N and Oryu S: 2004 A systematic study of light nuclei $2jA_j8$ using the AMD-method with realistic Paris AV14 nucleon-nucleon potentials, *Proc. of 17 International IUPAP Conf. on Few-Body Problems in Physics*, (2004) S198-S200.
- [49] Watanabe T: 2006 Calculation of a few nucleons systems by anti-symmetrized molecular dynamics. *Tokyo University of Science Doctoral thesis* (in Japanese)
- [50] Oryu S: 2012, Universal structure of the three-body system. *Phys. Rev.* **C86** 044001
- [51] Efimov V: 1970 Energy levels arising from resonant two-body forces in a three-body system. *Phys. Lett.* **B33** 563
- [52] Efimov V: 1973 Energy levels of three resonantly interacting particles. *Nucl. Phys.* **A210** 157
- [53] Oryu S, Hiratsuka Y and Watanabe T: 2016 Few-Body Problem in Nuclear Reactions: Beyond the horizon of the three-body Faddeev equations *EPJ Web of Conferences* **122** 08001 ; DOI: 10.1051/epjconf/201612208001 CNR*15
- [54] Hiratsuka Y, Oryu S and Watanabe T: 2014 $NN\pi$ Faddeev Calculation *talk in APFB6 Conference in Adelaide 2014*