

# H $\alpha$ images of nearby galaxy groups NGC193 and NGC940

S Hattings<sup>1</sup>, S I Loubser<sup>1</sup>, A Babul<sup>2</sup>, E O'Sullivan<sup>3</sup>

<sup>1</sup> Centre for Space Research, North-West University, Potchefstroom, South Africa

<sup>2</sup> University of Victoria, Canada

<sup>3</sup> Harvard-Smithsonian Centre for Astrophysics, USA

E-mail: sumarihatt@gmail.com

**Abstract.** A significant fraction of the baryons in galaxy groups and clusters is not locked up in stars and the interstellar medium (ISM), but exists in the form of extended multi-phase gas. The advantage of nearby galaxy groups is that this multi-phase gas can be studied via X-ray observations probing the hot gas ( $\approx 10^7 K$ ), and optical emission line observations (e.g. H $\alpha$  filaments) probing the ionised warm gas. Another advantage is that the effects of the galaxies' heating process of their surrounding gas (e.g. feedback from AGN) are more visible due to lower density and mass, and because groups contain fewer galaxies than rich clusters. Interaction between the different phases is important for galaxy formation and evolution due to the fact that at least 50% of galaxies can be found in groups and clusters.

The Complete Local-Volume Groups Sample (CLOGS) is an optically selected sample of 53 groups within 80 Mpc of which the X-ray observations (*Chandra / XMM-Newton*) have already been done and can be used to infer important physical properties, such as gas temperature and X-ray luminosity. In this project the H $\alpha$  images, observed using the WIYN 0.9m telescope on Kitt Peak, are analysed with the aim to compare the H $\alpha$  filaments to the X-ray emission images. Here we present preliminary results and a multi-wavelength comparison for two rich galaxy groups: NGC193 and NGC940. Despite their differences, we detect H $\alpha$  in the central members of both galaxy groups.

## 1. Introduction

In this study, we investigate the dominant early-type galaxy and surrounding members of two galaxy groups, NGC193, a galaxy group with its main member an X-ray bright radio galaxy with a powerful central jet, and the other group NGC940, with its main member an X-ray faint galaxy with only a weak radio point source. The two groups are part of the high-richness subsample of the Complete Local-volume Groups Sample (CLOGS). CLOGS is an optically-selected and statistically-complete sample of groups in the nearby Universe. Probing the luminous intra-group medium (IGM), CLOGS is specifically chosen and studied with optical observations for this project, since radio and X-ray bands [1] are already available. Important physical properties of the IGM are gas temperature and the total X-ray luminosity, both probing the environment in which the group members are located [2]. Together, the CLOGS observations can be used to investigate the role of active galactic nuclei (AGNs) in maintaining the thermal balance of the IGM, to name one example. The CLOGS galaxy groups were observed using the Giant Metrewave Radio Telescope (GMRT) in dual-frequency 235/610 MHz mode [1], with *Chandra / XMM-Newton*, and now also in H $\alpha$  with the WIYN 0.9m telescope.

The brightest, dominant early-type galaxies of groups are known for their typically high luminosity, and are old galaxies located near the centres of the group. These brightest group galaxies often reveal activity in their nuclei by radio emission & jets; depositing their energy back to the IGM. If conditions are favourable, the gas surrounding the brightest group member can cool down to form stars and feed the black hole. The latter results in AGN heating, which can heat the gas and cause the cycle to continue. In this article, we study two of the 53 groups, and we give a brief overview of their properties below.

### NGC0193<sup>1</sup>

This central dominant, jet-dominated, early-type galaxy, at the centre of the group NGC193, is a currently-active large-scale jet system located at a right ascension of 00h39m18.6s and a declination of +03d19m52s. Its jet components extend several tens of kpc away from the host galaxy, and are seen as very bright, straight jets. These radio jets contribute to the heating of the IGM by balancing the radiative energy losses and also maintaining the long-term thermal equilibrium [3].

The radio jets, known for their large cavities or “cocoon”-like structures, cause disturbances in the X-ray emitting gas [1]. These two bright, straight jets are revealed by the radio contours (see the right image of figure 2) and are in a low-surface brightness “cocoon”, extending perpendicular to the jet axis, out to a projected distance of  $\sim 30$  kpc from the centre - beyond the possible central cavity - and show very little correlation with the substructures detected in the X-ray image. Opposed to this, a correlation is observed between the outer border of the “cocoon” and the bright X-ray rim of the cavity in the northern region [4]. This may suggest that the cavity is possibly filled by the radio plasma in the diffuse “cocoon”. The X-ray bright rim represents a shell of gas, brightened on its edge, leaving little cool gas in the galaxy core.

In addition, this central member has X-ray emission-shells that surround its radio-lobes. The *Chandra* image (see the left image of figure 2), shows a bright ring with radius  $\sim 20$  kpc around the nucleus of the central X-ray peak. This bright ring is either the outer shell of a large single X-ray cavity projected towards the group centre, or the result of the superposition of two cavities along the line of sight [4].

### NGC0940<sup>2</sup>

This central dominant early-type galaxy in the centre of the group, NGC940, located at a right ascension of 02h29m27.5s and declination of +31d38m27.3s, shows vastly different properties to NGC0193. The radio emission shows a point source presenting a greater flux density at 610 MHz than at 235 MHz. No hot gas halo is detected in the X-rays [5].

The *XMM-Newton* image (see the left image of figure 5) shows a  $\sim 4$  kpc radius bright ring around the nucleus of the central X-ray peak, with the contours indicated in black.

Figure 5 (see the image on the right) shows the 610 MHz contours (indicated with green colours), overlaid on the smoothed optical Digital Sky Survey (DSS) image.

## 2. Observations

$H\alpha$  is a strong spectral line visible in the red part of the optical electromagnetic spectrum, which makes it easily traceable. This spectral line has a rest wavelength of  $6562.8\text{\AA}$  and is emitted when a hydrogen atom falls from its third ( $n = 3$ ) to second ( $n = 2$ ) lowest energy level.

In this project, the wide-field  $H\alpha$ -images used for both galaxy groups contain group members of different morphological types. One example of this is a spiral galaxy (refer to figure 3). This type is known for its spiral structure from its centre into the galactic disc and this morphological type makes up approximately 60% [6] of the local galaxy population in the current Universe. These galaxies contain more young, massive and bright stars in the spiral arms and therefore appear very bright.

Another type of galaxy is an elliptical galaxy, which has an elliptical shape with almost no features but a much more concentrated / compact structure. Such an example is shown in figure 3, the NGC0193 galaxy. The stars found in elliptical galaxies are older than the young bright stars found in the spiral arms of a spiral galaxy, and are known to be low-mass stars (opposed to the massive stars in spiral arms).

The  $H\alpha$ -filter,  $I_{H\alpha}$ , used to observe both galaxy groups for this project had a bandwidth of  $81.33\text{\AA}$  ( $k_{H\alpha}$ ) and a transmission rate of 90.58% ( $\tau_{H\alpha}$ ), whereas the r-band,  $I_r$ , had a bandwidth of  $1475.17\text{\AA}$  ( $k_r$ ) and a transmission rate of 92.83% ( $\tau_r$ ). To see the ionised gasses clearly, the r-band images must be used to subtract the continuum (i.e. stellar emission).

These images were scaled by using the formula [7]:

$$I_{[H\alpha]} = \frac{I_{H\alpha} \cdot k_r - I_r \cdot k_{H\alpha}}{\tau_{H\alpha} \cdot k_r - \tau_r \cdot k_{H\alpha}} \quad (1)$$

<sup>1</sup> Since the galaxy groups are named after their brightest early-type members, we denote the galaxy as NGC0193, and the group as NGC193.

<sup>2</sup> Similarly, we denote the other brightest early-type member as NGC0940, and the group as NGC940.


where the intensity  $I$  is in counts/sec.

This method then effectively shows the ionised hydrogen gasses very clearly, with prominent clumps of stars visible, especially in the spiral galaxy members of both galaxy groups. Such an example can be seen in figure 6 where the clumps of stars are clearly visible.

### 3. Discussion

#### 3.1. NGC193

Radio jets are known for its large cavities, heating the surrounding gas, which are often used to look at the power output of these jets [2]. The size of cavities are defined by matching an ellipse to their apparent shape. This is a very compact group since it has a higher density of galaxies in the centre (see Figure 1).




**Figure 1.** 15 Members of galaxy group NGC193 plotted according to their projected location in the group, colour-coded by brightness. The range in which each galaxy's magnitude lies, is indicated by the letter B.


Figure 1 corresponds to the coordinates of the members on the  $H\alpha$  image taken on the WIYN 0.9m telescope on Kitt Peak and the members were colour-coded according to their magnitudes where we show members with B-magnitude values between -18 and -21. Figure 1 shows that the main member, NGC0193 (number 7) is surrounded by five other bright members (purple indicated squares) of this galaxy group.

X-ray and radio data for this group's main member, NGC0193, indicate full- and low-resolution contours on the Chandra and optical image, respectively (see Figure 2). The two very bright jet components are revealed by radio contours.

For three members of this group, NGC0193, NGC0198 and NGC0200, the r-band was subtracted from the continuum and scaled (see Figure 3) by a formula [7] discussed in the previous section. The ionised hydrogen gasses are shown very clearly for these three members with prominent clumps of stars for NGC0198 and NGC0200 (images on the right).




**Figure 2.** X-ray and radio data for group NGC193. *Left:* GMRT 610 MHz full-resolution contours overlaid on the smoothed, 0.3-2.0 keV Chandra image. *Right:* GMRT 235 MHz low-resolution contours on the optical image. Both images' radio contours are spaced by a factor of 2 [4]



**Figure 3.** Three bright members NGC0193, NGC0198 and NGC0200 (in order from top to bottom) of the galaxy group NGC193. H $\alpha$  filter images (left), r-band images (middle) and the r-band subtracted from the H $\alpha$  (right).

### 3.2. NGC940


For this galaxy group, no diffuse hot halo has been detected in X-ray observations, but rather a point-source. This galaxy group is much less compact in terms of galaxies, since it only has nine members with the main member (see Figure 6) one of the brightest galaxies in this group, which was expected.




**Figure 4.** Nine members of galaxy group NGC940 plotted according to their projected location in the group, colour-coded by brightness. The range in which each galaxy's magnitude lies, is indicated by the letter B.

Figure 4 indicates that the main member, NGC0940 (number 5) is surrounded by one other bright member (purple indicated squares) of this galaxy group. The members were colour-coded according to their B-magnitude values between -18 and -21.

X-ray and radio data for main member NGC0940 reveal low- and full-resolution contours on the *XMM-Newton* and optical DSS image, respectively (see Figure 5). The contours overlaid on both images indicates no jet components, since this main member does not have a hot halo but is rather detected as a point source.



**Figure 5.** X-ray and radio data for group NGC940. *Left:* GMRT 235 MHz low-resolution contours in black overlaid on the adaptively smoothed, 0.3-2.0 KeV of *XMM-Newton* image. *Right:* GMRT 610 MHz full-resolution contours in green overlaid on the optical DSS image. [1]



**Figure 6.** Three bright members, NGC0940, NGC0931 and UGC01963 (in order from top to bottom) of the galaxy group NGC940.  $H\alpha$  filter images (left), r-band images (middle) and the r-band subtracted from the continuum (right).

Figure 6 shows that three bright members were scaled by making use of the formula [7] as discussed in the previous section. After the r-band of each member was subtracted from the continuum (images on the right), the ionised hydrogen gases were clearly visible, showing the prominent clumps of stars in members NGC0931 and UGC01963.

#### 4. Conclusions

In this study we have investigated surrounding members of two rich galaxy groups, NGC193 and NGC940 which are part of the CLoGS sample and can be used to investigate the role of AGNs to maintain the thermal balance of the IGM. Here we present preliminary results for both galaxy groups, detecting  $H\alpha$  in their central members - despite of the two galaxies' different properties from X-ray and radio observations (e.g. jet-dominated vs point-source). Therefore, the environment and the IGM surrounding their dominant central galaxies are very diverse. Yet, intriguingly,  $H\alpha$ -emission is detected in both dominant galaxies, and the physical connection between the different phases (e.g. hot gas, and warm ionised gas) is not obvious. It will be interesting to now investigate the other members of the galaxy groups (surrounding the dominant member), as well as extend the study to include more CLoGS groups to thoroughly explain the origin of the ionised  $H\alpha$  gas.

#### Acknowledgments

Acknowledgements SH is supported by the National Astrophysics and Space Science Program (NASSP). SIL is aided by a Henri Chrétien International Research Grant administered by the American Astronomical Society and the National Research Foundation (NRF).

#### References

- [1] Kolokythas K et al 2018 The complete local-volume groups sample-ii: a study of the central radio galaxies in the high-richness sub-sample *MNRAS* **481** 1550
- [2] O'Sullivan E et al 2017 The complete local volume groups sample-i: sample selection and X-ray properties of the high-richness subsample *MNRAS* **472** 1482
- [3] Babul A et al 2002 Physical implications of the X-ray properties of galaxy groups and clusters *MNRAS* **330** 329
- [4] Giacintucci S et al 2011 A combined low-radio frequency / X-ray study of galaxy groups i: giant metrowave radio telescope observations at 253 MHz and 610 MHz *ApJ* **732** 95
- [5] Tully R B 2015 Galaxy groups: a 2-mass catalog *ApJ* **1499** 171
- [6] Loveday J 1996 The APM bright galaxy catalogue *MNRAS* **278** 1025
- [7] Terlevich R, Melnick J, Masegosa J, Moles M, Copetti M V F 1991 Spectrophotometric catalogue of HII galaxies *A&AS* **91** 285