



Contribution ID: 47

Type: Oral Presentation

Light flavor symmetry breaking for heavy baryons

Friday, 13 July 2012 08:20 (20 minutes)

Abstract content
 (Max 300 words)

We are interested in the soliton description of baryons with a single heavy quark (charm or bottom). In this approach such baryons emerge as composites of a soliton of mesons built from light quarks (up, down, strange) and a meson field that is bound to the soliton and contains a heavy quark. The soliton must then be quantized as a diquark because the fermionic character arises from binding to the heavy meson field. We are particularly interested in baryons that contain strange flavors; in the quark model that corresponds to, say, up-strange-bottom. Thus the flavor symmetry breaking among the light quarks must be incorporated when constructing diquark states. Here we present computations of the diquark eigen-energies and eigen-functions that incorporates all orders of the light flavor symmetry breaking. We also compare these results to a leading order treatment of flavor symmetry breaking. This is a first step towards a comprehensive description of heavy baryons in a soliton model.

Apply to be
 consider for a student
 award (Yes / No)?

Yes

Level for award

- (Hons, MSc,

- PhD)?

PhD

Main supervisor (name and email)
 -br>and his / her institution

Prof. Herbert Weigel, weigel@sun.ac.za Stellenbosch University

Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?

No

Primary author: Mr BLANCKENBERG, Jaco (Stellenbosch University)

Co-author: Prof. WEIGEL, Herbert (Stellenbosch University)

Presenter: Mr BLANCKENBERG, Jaco (Stellenbosch University)

Session Classification: Theoretical

 $\textbf{Track Classification:} \ \ \mathsf{Track} \ \mathsf{G} \ \mathsf{-} \ \mathsf{Theoretical} \ \mathsf{and} \ \mathsf{Computational} \ \mathsf{Physics}$