

# Decoherence-assisted transport in quantum networks

**Adriana Marais, Ilya Sinayskiy and Francesco Petruccione.**

Quantum Research Group and National Institute for Theoretical Physics, School of Chemistry and Physics, University of KwaZulu-Natal, Durban, 4001, South Africa

E-mail: [adrianamarais@gmail.com](mailto:adrianamarais@gmail.com), [sinayskiy@ukzn.ac.za](mailto:sinayskiy@ukzn.ac.za), [petruccione@ukzn.ac.za](mailto:petruccione@ukzn.ac.za)

**Abstract.** The dynamics of a quantum network under the influence of decoherence are studied. This work is a generalisation of previous research on decoherence-assisted transport in a dimer system [I. Sinayskiy, A. Marais, F. Petruccione and A. Ekert, *Phys. Rev. Lett.* 108, 020602 (2012)]. The model under investigation consists of a homogenous fully connected quantum network in contact with an environment of spins. Exact analytical expressions for the transition probabilities are obtained. It is shown that there exist well-defined ranges of parameters for which decoherent interaction with the environment assists energy transfer in the quantum network. This model of decoherence-assisted energy transfer is applied to energy transfer in the Fenna-Matthews-Olson complex.

## 1. Introduction

Recently, evidence of quantum coherence has been detected in biological systems at physiological temperature, including the photosynthetic light-harvesting complexes of a species of green sulphur bacteria [1] and two species of marine cryptophyte algae [2]; organisms well-adapted to photosynthesise under low-light conditions. Light-harvesting complexes act as antennas, absorbing photons and transferring the resulting excitation energy through a network of photoactive pigments held in well-defined orientations and configurations by a scaffold of proteins, to the reaction centre, where the secondary photosynthetic process of charge separation takes place. The electronic excitation energy transfer (EET) happens on a scale of picoseconds and with a quantum efficiency of over 95% [3]. The surprising phenomenon of quantum coherence in warm, noisy, complex and yet remarkably efficient energy transfer systems has led to discussion about the role of the protein environment in the energy transfer process and the degree to which it may contribute to this efficiency [4].

Modelling the complexity of the environment is a challenge. The protein-solvent environment interacts strongly with the pigments due to its polarity and as a result can have a significant effect on the quantum dynamics, which will therefore in general be non Markovian [5]. Such non-Markovian effects have widely been taken into account [5, 6], but so far, all have been within spin-boson models of excitons within a protein medium. While any biological system is always in contact with a bosonic environment, interaction with a more structured environment such as a spin bath is more likely to assist quantum efficiency and also induces intrinsically non-Markovian dynamics [7].

With the aim of exact solvability, we investigate the relationship between environmental spin

degrees of freedom and the efficiency of the EET process described by the Hamiltonian  $H_{ex}$

$$H_{ex} = \sum_j E_j |j\rangle\langle j| + \sum_{i \neq j} V_{ij} |i\rangle\langle j|, \quad (1)$$

where the site energies of the pigments are given by  $E_j$ , and the EET couplings by  $V_{ij}$ . We review the case of the dimer system; generalise this model to a fully connected network; and finally apply this model to the Fenna-Matthews-Olson (FMO) antenna complex [10].

## 2. Dimer

For a dimer with Hamiltonian  $H_d = \varepsilon_1 |1\rangle\langle 1| + \varepsilon_2 |2\rangle\langle 2| + J |1\rangle\langle 2| + |2\rangle\langle 1|$ , the maximum transfer probability for a single excitation  $\text{Max}[P_{1 \rightarrow 2}(t)]$  is given by  $J^2/(J^2 + \Delta^2)$  where  $J$  is the amplitude of transition, and the detuning  $\Delta$  is given in terms of the energy levels of the dimer as  $(\varepsilon_2 - \varepsilon_1)/2$ . Certain transfer is achieved when  $\Delta = 0$  at time  $t = \pi/(2J)$ , or when there is resonance between the energy levels in the system.

In a recent article [8], it is shown that there exist well-defined ranges of parameters for which a purely decoherent interaction with environmental spins in a spin star configuration [7] assists energy transfer in the dimer system. For a dimer with each level coupled to a spin bath at zero temperature, the Hamiltonian of the total system is given by

$$H = H_d + H_B + H_I. \quad (2)$$

Each environment  $B_j$  consists of  $n_j$  spin-half particles

$$H_B = \sum_{j=1}^2 H_{B_j} = \alpha_j \sum_{j=1}^2 \sum_{k=1}^{n_j} \frac{\sigma_z^{k,j}}{2}, \quad (3)$$

where  $\sigma_z$  are the Pauli matrices. The purely decoherent interaction between each site  $j$  in the system and the corresponding spin bath is modelled by

$$H_I = \sum_{j=1}^2 H_{I_j} = \sum_{j=1}^2 \sum_{k=1}^{n_j} \gamma_j |j\rangle\langle j| \frac{\sigma_z^{k,j}}{2}. \quad (4)$$

The Hamiltonian of the environment  $H_B$  commutes with the Hamiltonian of interaction  $H_I$  and therefore the state of the total system is always in a product state of the network and the baths. As a result, the effective Hamiltonian for the total system with spin baths at zero temperature is given by

$$H = \sum_{j=1}^2 \varepsilon_j |j\rangle\langle j| + \sum_{\substack{i,j=1 \\ i \neq j}}^N J |i\rangle\langle j|, \quad (5)$$

where  $\varepsilon_j = \varepsilon - \gamma_j n_j/2$ .

For the Hamiltonian  $H$ , the maximum transfer probability  $\text{Max}[P_{1 \rightarrow 2}(t)]$  is given by  $J^2/(J^2 + \Delta'^2)$  where in this case the detuning is given by  $\Delta' = (\varepsilon'_2 - \varepsilon'_1)/2$ , and  $\varepsilon'_j = \varepsilon_j - n_j \gamma_j/2$  are the shifted energy levels of the dimer as a result of coupling with strength  $\gamma_j$  with each of the spin baths with number of spins  $n_j$ . Certain transfer is similarly achieved when  $\Delta' = 0$ , which in this case is possible for a wide range of the parameters  $\gamma_j$  and  $n_j$ .

This effect persists at physiological temperature, where transfer probabilities of nearly 90% can be achieved in the dimer at 300K for biologically relevant parameters [8].

### 3. Fully-connected quantum network

For a fully connected network of  $N$  qubits interacting via homogeneous Heisenberg  $XX$  coupling with coupling strength  $2J$  and with equal site energies  $\varepsilon$ , the effective Hamiltonian in the single excitation subspace is given by

$$H_N = \sum_{i=1}^N \varepsilon |i\rangle\langle i| + \sum_{\substack{i,j=1 \\ i \neq j}}^N J |i\rangle\langle j|. \quad (6)$$

By coupling  $N - k$  of the sites in the fully connected network to independent spin environments in symmetric star configurations, the Hamiltonian of the total system is given by

$$H_k = H_N + H_B + H_I, \quad (7)$$

with  $H_B$  and  $H_I$  defined as previously, but with  $j = 1, \dots, N - k$ . We can then write the effective Hamiltonian for the total system, with the baths arbitrarily coupled to the last  $N - k$  sites, as

$$H_k = \sum_{j=1}^k \varepsilon |j\rangle\langle j| + \sum_{j=k+1}^N \varepsilon_j |j\rangle\langle j| + \sum_{\substack{i,j=1 \\ i \neq j}}^N J |i\rangle\langle j|, \quad (8)$$

where  $\varepsilon_j = \varepsilon - \gamma_j n_j / 2$ .

In the case where  $k = N$  and all levels have equal energy  $\varepsilon$ , the Hamiltonian  $H_N$  is given by Eq. (1). In this case the maximum probability of purely coherent transfer through the network is

$$\text{Max}[P_{I \rightarrow F}(t)] = \frac{4}{N^2}, \quad (9)$$

at time  $t = \pi / (NJ)$ . Our study will focus on whether decoherent interaction between a fully-connected network and environmental spins can enhance energy transport through the network.

We analyse a range of cases where spin baths are added to the initial, final and intermediate network sites, and find that decoherent interaction with the spin baths in general increases transfer probability through the network, and furthermore that these effects persist at physiological temperature [9].


As an example, we consider here the case of coupling both the initial and final sites to spin baths, with Hamiltonian  $H_{N-2}$  given by Eq. (8) with  $k = N - 2$ . In this case, it can be shown that when the shifted energy levels are equal,  $\varepsilon_1 = \varepsilon_2$ , there exist times for which the transfer probability is arbitrarily close to 1.

We now show that such an effect persists at physiological temperatures by considering the Hamiltonian  $H_{N-2}$  with spin baths at a temperature of 300K coupled to the initial and final sites.

In this case the initial state of the bath is given by the canonical distribution

$$\rho_B(0) = \prod_{i=1}^2 \frac{1}{Z_i} e^{-\beta \alpha_i S_i^z}, \quad (10)$$

where  $Z_i$  is the partition function of the corresponding spin bath.




**Figure 1.** Graph of the maximum of the probability of transition  $\text{Max}[P_{I \rightarrow F}(t)]$  at 300K in a 10-site fully connected homogeneous network, with isolated site energies  $\varepsilon = 100\text{ps}^{-1}$ , and spin baths coupled to each of the initial and final sites, with  $n_1 = n_2 = 10$  and bath energy parameter  $\alpha_1 = \alpha_2 = 100\text{ps}^{-1}$ .

In Fig. 1 the maximum of the probability of transition  $\text{Max}[P_{I \rightarrow F}(t)]$  for such a system is plotted as a function of the coupling constants  $\gamma_1$  and  $\gamma_2$  with spin baths coupled to each of the initial and final sites. It can be seen that in regions where  $\varepsilon_1 \approx \varepsilon_2 \neq \varepsilon$ , transfer probabilities of up to 72 % are achieved.

#### 4. FMO complex

The first evidence of quantum coherence in photosynthetic antennas at physiological temperature was detected green sulphur bacteria and cryptophyte algae [1, 2], both organisms able to photosynthesise efficiently at low light intensities. Green sulphur bacteria uniquely contain a complex called the Fenna-Matthews-Olson (FMO) complex [10], which mediates excitation energy transfer from the antenna to the reaction centre [11]. The site energies and optical transition energies for the FMO complex of *Chlorobium tepidum* used here were calculated by Adolphs and Renger [12]. Quantum coherent EET through the bare excitonic system without adding environmental contributions to the Hamiltonian  $H_{ex}^{FMO}$  happens with a low probability: for the transfer of the excitation to from site 1 to site 3, the probability of transfer is just 4.2%. We now investigate the effect of decoherent interaction with environmental spins at 300K on the process of EET in the FMO complex.

With the spin baths at a temperature of 300K, we calculate the maximum probability of transfer during the first picosecond, for equal environmental couplings  $\gamma$  between 0 and  $250\text{ps}^{-1}$  at each site. We find that for a total of 10 spins, distributed between the seven network sites as follows:  $n_1 = 2, n_4 = 8$  and  $n_{2,3,5,6,7} = 0$ , and with spin bath energy constant  $\alpha = 150\text{ps}^{-1}$ , the maximum transfer probability is 86% (see Fig. 2). This is a vast increase in the transfer probability from the case with no spin baths, where the probability is 4.3%. Therefore, in this case decoherence



**Figure 2.** The probability of transfer  $P_{1\rightarrow 3}(\gamma, t)$  for the FMO complex with sites coupled to a spin baths at 300K with numbers of spins at each site  $n_1 = 2, n_4 = 8$  and  $n_{2,3,5,6,7} = 0$  and spin bath energy constant  $\alpha=150\text{ps}^{-1}$ . The maximum transfer probability is 86%.

assists the efficiency of quantum coherent EET in the FMO complex.

## 5. Conclusion

The recent detection of quantum coherence in biological systems that are remarkably efficient in transferring excitation energy at physiological temperatures, has led to investigation as to whether this coherence contributes to the efficiency. Here, we have investigated the influence of environmental spins on quantum coherent transfer. We have shown through the derivation of analytical expressions that the transfer probabilities through a fully connected quantum network are improved as a result of decoherent interaction with environmental spins, and that in some cases certain transfer can be achieved. Moreover, this effect is shown to persist at physiological temperatures. We apply this model to the FMO complex, and find that coupling the network sites with environmental spins at physiological temperature improves transport through the network for the considered case. These promising results motivate further study of biological transport systems where environmental spins may play an important role in the dynamics.

## Acknowledgments

The authors would like to thank Artur Ekert and Alastair Kay for valuable and interesting discussions. This work is based upon research supported by the South African Research Chair Initiative of the Department of Science and Technology and National Research Foundation.

## References

- [1] G. Panitchayangkoon, D. Hayes, K. A. Fransted, J. R. Caram, E. Harel, J. Wen, R. E. Blankenship and G. S. Engel., Proc. Natl. Acad. Sci. U. S. A. 107, 12766 (2010).
- [2] E. Collini, C.Y. Wong, K.E. Wilk, P.M G. Curmi, P. Brumer and G.D. Scholes, Nature 463, 08811 (2010).
- [3] B. R. Green and W. W. Parson (Eds.), Light-harvesting Antennas in Photosynthesis (Kluwer Academic Press, Dordrecht, 2003).
- [4] M. Mohseni et al., J. Chem. Phys. 129, 174106 (2008); M.B. Plenio and S.F. Huelga, New J. Phys. 10, 113019 (2008); K. Furuya, and G.J. Milburn, New J. Phys. 12, 083033 (2010); S. Lloyd and M. Mohseni, New J.

- Phys. 12, 075020 (2010); M. Sarovar et al., Nature Physics 6, 462 (2010); F. Caruso et al., Phys. Rev. A 81, 062346 (2010); I. Kassal and A. Aspuru-Guzik, arXiv:1201.5202 (2012).
- [5] A. Ishizaki and G. R. Fleming, J. Chem. Phys. 130, 234110 (2009); 130, 234111 (2009).
  - [6] V. I. Novoderezhkin and R. van Grondelle, Phys. Chem. Chem. Phys. 12, 7352 (2010).
  - [7] H.-P. Breuer, D. Burgarth, and F. Petruccione, Phys. Rev. B 70, 045323 (2004).
  - [8] I. Sinayskiy, A. Marais, F. Petruccione and A. Ekert, Phys. Rev. Lett. 108, 020602 (2012).
  - [9] A. Marais, I. Sinayskiy, F. Petruccione and A. Ekert (in preparation).
  - [10] R. E. Fenna and B. W. Matthews, Nature 258, 573 (1975).
  - [11] R. Blankenship, J. Olson and M. Miller, A.I.P.H. 2, 399 (2004).
  - [12] J. Adolphs and T. Renger, Biophys. J. 91, 2778 (2006).