

Contribution ID: 451

Type: Oral Presentation

Fermi edge singularity and finite frequency spectral features in a semi-infinite 1D wire

Friday, 13 July 2012 11:00 (20 minutes)

Abstract content
 (Max 300 words)

We theoretically study a charge qubit interacting with electrons in a semi-infinite 1D wire. The system displays the physics of the Fermi edge singularity, i.e. a singularity in the qubit tunneling rate due to a proliferation of long wavelength excitations in the wire. Our results generalize known results for the Fermi-edge system to the regime where excitations induced by the qubit can resolve the spatial structure of the scattering region. We find resonant features in the qubit tunneling rate as a function of the qubit level splitting. They occur at integer multiples of Planck's constant times the Fermi velocity divided by l. Here l is the distance from the tip of the wire to the point where it interacts with the qubit. These features are due to a single coherent charge fluctuation in the electron gas, with a half-wavelength that fits into l an integer number of times. As the coupling between the qubit and the wire is increased, the resonances are washed out. This is a clear signature of the increasingly violent Fermi-sea shake-up that accompanies strong coupling.

Apply to be
 consider for a student
 award (Yes / No)?

No

Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?

No

Primary author:Dr SNYMAN, Izak (NITheP)Co-author:Dr SHEIKHAN, Ameneh (NITheP)Presenter:Dr SNYMAN, Izak (NITheP)Session Classification:Theoretical

Track Classification: Track G - Theoretical and Computational Physics