



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA

Contribution ID: 387

Type: **Poster Presentation**

Enhanced luminescence from Tb for the mixed spinel $\text{Mg}_{1-x}\text{Zn}_x\text{Al}_2\text{O}_4$

Thursday, 12 July 2012 17:30 (2 hours)

Abstract content (Max 300 words)

Spinel, which have the general formula AB_2O_4 , often occur naturally as minerals but are also synthesized and studied for their interesting electrical, magnetic and optical properties. MgAl_2O_4 finds diverse applications due to its mechanical strength, chemical inertness, relatively low density, high melting point, high thermal shock resistance, low thermal expansion coefficient, resistance to neutron irradiation and low dielectric loss. It has also been used as a phosphor host activated by a variety of transition metal and lanthanide ions. ZnAl_2O_4 is widely used as a catalyst and has recently attracted much attention as a phosphor host. Both have been studied separately as possible hosts for Tb, but in this work they were compared directly. Results were also obtained for Tb-doped mixed spinels $\text{Mg}_x\text{Zn}_{1-x}\text{Al}_2\text{O}_4$. Although the lattice constant changes little with composition, the bandgap of MgAl_2O_4 (7.8 eV) is double that of ZnAl_2O_4 (3.9 eV). Nanocrystalline powder samples with a particle size of about 25 nm were prepared using the combustion method. For $\text{MgAl}_2\text{O}_4\text{:Tb}(0.5 \text{ mol}\%)$ both green emissions due to $5\text{D}_4\text{-}7\text{F}_J$ transitions (with the most intense $5\text{D}_4\text{-}7\text{F}_5$ transition at 544 nm) and blue emissions due to $5\text{D}_3\text{-}7\text{F}_J$ transitions were observed. Less intense green emissions were observed for $\text{ZnAl}_2\text{O}_4\text{:Tb}(0.5 \text{ mol}\%)$ and no blue emission occurred. Both samples had similar excitation spectra (for 544 nm emission), with a peak near 230 nm which is attributed to the Tb $4\text{f-}5\text{d}$ transition. This was unexpected due to the large difference in their bandgaps. This excitation wavelength corresponds to an energy of 5.4 eV which is higher than the bandgap of ZnAl_2O_4 . Therefore a fraction of the incident light will be absorbed by this host and not be available to excite the Tb ions, which corresponds to the observation of poorer luminescence from the $\text{ZnAl}_2\text{O}_4\text{:Tb}$. The absence of blue emission peaks is usually attributed concentration quenching, but since the same Tb concentration was used for the $\text{MgAl}_2\text{O}_4\text{:Tb}$ where blue emissions did occur, it is rather suggested that because of the smaller bandgap of ZnAl_2O_4 , the 5D_3 level lies close to or inside the conduction band and this prevents transitions from this level. The maximum green emission was measured for the mixed spinel $\text{Mg}_{0.75}\text{Zn}_{0.25}\text{Al}_2\text{O}_4\text{:Tb}(0.5 \text{ mol}\%)$, although the blue emissions from this sample were less than for the MgAl_2O_4 host.

Apply to be considered for a student award (Yes / No)?

Yes

Level for award (Hons, MSc, PhD)?

PhD

Main supervisor (name and email) and his / her institution

**Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?**

Yes

Primary author: Mr TABAZA, Wael (Physics Dept, UFS)

Co-authors: Prof. SWART, Hendrik (Physics Dept, UFS); Dr KROON, Ted (Physics Dept, UFS)

Presenter: Mr TABAZA, Wael (Physics Dept, UFS)

Session Classification: Poster Session

Track Classification: Track A - Division for Condensed Matter Physics and Materials