SAIP2019



Contribution ID: 110

Type: Oral Presentation

Top-quark background estimation for physics Beyond Standard Model in the dilepton and jets final state with the ATLAS detector

Thursday, 11 July 2019 10:00 (20 minutes)

This analysis presents the top-quark background estimation for the $H \to Sh$ channel using the full Run-2 data, corresponding to an integrated luminosity of 139 fb⁻¹ of pp collisions at $\sqrt{s} =$ 13 TeV. The final state for this analysis is formed by two oppositely-charged leptons ($e^{\pm}\mu^{\mp}, \mu^{\pm}e^{\mp}, e^{\pm}\mu^{\mp} + \mu^{\pm}e^{\mp}$) and jets. In this analysis the dominant Standard Model background are the top-quark processes ($t\bar{t}$ and Wt). The top control/validation regions are defined in the 1 b-jet and 2 b-jets phase space, respectively. The construction of the top-quark validation region is defined to correct the Monte Carlo mismodeling observed in the leptonic kinematics. Finally, the derived normalization factor in the top-quark control region is applied in the top-quark processes in both the control region and the signal region.

Apply to be
 considered for a student
 award (Yes / No)?

Yes

Level for award
 (Hons, MSc,
 PhD, N/A)?

MSc

Primary author: Mr MONNAKGOTLA, Jeremiah (University of the Witwatersrand)
Co-author: Prof. MELLADO, Bruce (University of the Witwatersrand)
Presenter: Mr MONNAKGOTLA, Jeremiah (University of the Witwatersrand)
Session Classification: Nuclear, Particle and Radiation Physics

Track Classification: Track B - Nuclear, Particle and Radiation Physics