



Contribution ID: 242

Type: Oral Presentation

## Bow shocks formed by massive runaway stars in 3D

Thursday, 29 July 2021 12:30 (15 minutes)

Hyper-runaways are stars moving at supersonic speeds through the interstellar medium; they can be thought of as a subset of runaway stars but moving with velocities that are comparable to the Galactic escape velocity ( $\sim 500$  km/s). Because of the strong stellar winds and high space velocities, we expect massive (hyper)runaway stars to produce bow shocks. We use PLUTO, a magneto-hydrodynamics grid code, to simulate these bow shocks, performing axi-symmetric hydrodynamic simulations in 3-dimensions while including thermal conduction and detailed radiative cooling processes. In this talk we will present our results for a range of stellar velocities ( $100 \leq v_{\text{star}} \leq 500$  km/s) and discuss the implications for potentially observing hyper-runaways.

**Apply to be considered for a student ; award (Yes / No)?**

Yes

**Level for award;(Hons, MSc, PhD, N/A)?**

MSc

**Primary author:** RAMALATSWA, Katlego (University of Cape Town)

**Co-author:** Prof. MOHAMED, Shazrene (SAAO)

**Presenter:** RAMALATSWA, Katlego (University of Cape Town)

**Session Classification:** Astrophysics

**Track Classification:** Track D1 - Astrophysics