

# Photosynthesis Underneath Stones in the Namib Desert

**Michal Gwizdala<sup>1,2</sup>, Pedro H. Lebre<sup>3</sup>, Don A. Cowan<sup>3</sup>, Tjaart P.J. Krüger<sup>1,2</sup>**

<sup>1</sup>*Cnr Lynnwood and University Roads Hatfield Campus, Pretoria 0002, South Africa. Department of Physics, University of Pretoria;*

<sup>2</sup>*Cnr Lynnwood and University Roads Hatfield Campus, Pretoria 0002, South Africa. Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria;*

<sup>3</sup>*Cnr Lynnwood and University Roads Hatfield Campus, Pretoria 0002, South Africa. Centre for Microbial Ecology and Genomics, Department of Biochemistry, Genetics and Microbiology, University of Pretoria.*

*Corresponding author e-mail address: gwizdala.michal@gmail.com*

## 1. Introduction

In hyper-arid soil environments, photosynthetic microorganisms are largely restricted to hypolithic habitats. They occupy the ventral surfaces of translucent pebbles in desert pavements.

## 2. Results

We combined fluorometric, spectroscopic, biochemical and metagenomic approaches to investigate the light transmission properties of quartz stones in the Namib Desert, and assess the photosynthetic activity of the underlying hypolithic cyanobacterial biofilms. Quartz pebbles greatly reduced the total photon flux to the ventral surface biofilms and filtered out primarily the short wavelength portion of the solar spectrum. Chlorophylls d and f were not detected in biofilm pigment extracts; however, hypolithic cyanobacterial communities showed some other evidence of adaptation to sub-lithic conditions, like the prevalence of genes encoding Helical Carotenoid Proteins, which are associated with desiccation stress. Under water-saturated conditions, hypolithic communities showed no evidence of light stress, even when the quartz stones were exposed to full midday sunlight. This work adds to an understanding of the mechanisms behind the unique robustness of photoautotrophic organisms in extreme environments.