



Contribution ID: 192

Type: not specified

Investigation into the geological radiation levels and evaluation of hazard parameters in soil and rock specimens taken from mining sites across North-Eastern Nigeria.

Monday, 18 November 2024 16:45 (15 minutes)

Investigation into the geological radiation levels and evaluation of hazard parameters in soil and rock specimens taken from mining sites across North-Eastern Nigeria.

Michael C. Ohakwere-Eze¹, Musa Nafiu², Shiv Kumar Singh³, Momoh Kabiru¹, and John Simon¹

¹Department of Physics, Faculty of Science, National Open University of Nigeria.

²Department of Physics, Federal University of Kashere, Gombe State.

³Department of Physics, Federal University of Transportation Daura, Katsina State.

Corresponding author e-mail address: michael.ohakwereze@gmail.com

1. Introduction

This research quantitatively evaluates the natural radioactivity levels and geological radiation hazard parameters of soil and rock specimens obtained from mining locations in North-Eastern Nigeria, by means of gamma-ray spectroscopy. There have been potential public health risks associated with the use of soil and rock from mining locations in North-Eastern Nigeria, specifically the study areas. A total of twenty-eight samples were systematically gathered from Nahuta and Kashere locations. Through gamma spectrometry employing a NaI (TI) detector, the natural radioactivity levels of ²³⁸U, ²³²Th and ⁴⁰K were determined for each and every sample. The findings indicated that the mean activity concentrations of ²²⁶Ra, ²³²Th, and ⁴⁰K in Nahuta are 46.13±4.78 Bq/Kg, 34.10±3.02 Bq/Kg and 473.94±5.41 Bq/Kg for the soil samples respectively, and 32.91±0.49 Bq/Kg, 40.70±0.41 Bq/Kg, and 578.18±4.28 Bq/Kg for the rock samples respectively. The corresponding mean activity concentrations of ²²⁶Ra, ²³²Th, and ⁴⁰K in Kashere are 17.99±4.18 Bq/Kg, 23.73±1.78 Bq/Kg, and 191.65±3.15 Bq/Kg, for the soil samples, and 20.24±3.72 Bq/Kg, 29.09±1.78 Bq/Kg, and 148.36±3.15 Bq/Kg, for the rock samples respectively. An analysis of radiation risk parameters (D, AEDE, Raeq, Hex, Hin, AGDE and ELCR) has been conducted and findings explored. While the samples from the Kashere region fall within the international recommended levels, elevated readings of certain radiation health parameters are observed in the Nahuta region, posing serious public health risk due to utilization of the soil and rock from this area in construction activities.

1. Results

Larger activity concentration values for both soil and rock samples are found in Nahuta when compared with same for Kashere (Figure 1). In Nahuta, the ranges of radionuclide concentration for ²²⁶Ra, ²³²Th, and ⁴⁰K are 16.20±4.26 to 53.94±4.72 Bq/Kg with an average 46.13±4.78 Bq/Kg, 13.02±3.32 to 62.71±4.37 with an average 34.10±3.02 Bq/Kg and 76.4±8.44 to 1126.53±8.90 Bq/Kg with an average 473.94±5.41 Bq/Kg, respectively. Conversely, the radionuclide concentration of ²²⁶Ra, ²³²Th, and ⁴⁰K in Kashere ranges from 17.99±4.18 to 29.56±8.91 Bq/Kg with an average 17.99±4.18 Bq/Kg, 14.60±0.47 to 34.60±0.71 Bq/Kg with an average 23.73±1.78 Bq/Kg, and 85.74±6.03 to 316.00±2.99 Bq/Kg with an average 191.65±3.15 Bq/Kg, respectively. The activity concentration in rock samples is higher in Nahuta, with values for ²²⁶Ra, ²³²Th, and ⁴⁰K ranging from 9.74±1.26 to 92.78±0.28 Bq/Kg with an average 32.91±0.49 Bq/Kg, 14.33±0.26 to 71.04±0.10 Bq/Kg with an average 40.70±0.41, and 36.67±6.05 to 1136.50±5.54 Bq/Kg with an average 578.18±4.28, respectively. In Kashere, the ranges are 11.07±8.91 to 38.06±0.17 with an average 20.24±3.72 Bq/Kg for ²²⁶Ra, 13.47±6.23 to 42.43±0.36 Bq/Kg with an average 29.09±1.78 Bq/Kg for ²³²Th, and 13.77±2.99 to 326.89±2.33 Bq/Kg with an average 148.36±3.15 Bq/Kg for ⁴⁰K.

The concentrations of radionuclides ²²⁶Ra, ²³²Th, and ⁴⁰K in soil and rock samples, as reported by [1], average

at 32, 45, and 420 Bq/kg, respectively, worldwide. Analysis indicates that the average radionuclide concentration of ^{226}Ra , ^{232}Th , and ^{40}K in soil and rock samples from Kashere region are lower than the global averages, suggesting no imminent radiological threat to the local population. Conversely, the average activity concentration of ^{226}Ra and ^{40}K in soil and rock samples from Nahuta exceeds worldwide averages. These elevated levels are attributed to mining activities in the study area, which is abundant in natural radionuclides.

The calculated mean activity concentration of ^{226}Ra is higher in the soil sample collected from Nahuta compared to Kashere. Nevertheless, compared to Nahuta, Kashere has slightly elevated levels in samples derived from rocks. This implies that the geological formation and mining exploration in Nahuta may have resulted in higher concentrations of radium than in Kashere where the rocks may contain higher concentrations of radium. The mean activity concentrations of ^{232}Th in the soil and rock samples collected from Nahuta are higher than those obtained for Kashere. The higher thorium levels in Nahuta could be explained by the kind of rocks that compose the formations present in the area. The mean activity concentrations of ^{40}K are higher in the soil and rock samples collected from Nahuta as compared to Kashere. Potassium-40 being present in many rocks occurs at higher concentrations in Nahuta, signifying a higher concentration of potassium bearing minerals in the area.

As observed in Figure 1, there are higher concentrations of ^{226}Ra , ^{232}Th , and ^{40}K , particularly in the soil samples obtained from Nahuta, which may pose a radiological health hazard to the inhabitants of the area. This is a health issue because prolonged exposure to these radionuclides contributes to cancer risk and other illnesses. Also, the sampled higher concentrations of these radionuclides suggest the need for environment management and monitoring systems, which should mitigate the impacts of mining and other agricultural activities. The results emphasize the necessity to follow the legal requirements concerning the radiation safety in the region with anthropogenic activities.

Fig. 1. Variation of the mean activity concentrations of ^{226}Ra , ^{232}Th , and ^{40}K given in Bq/Kg for Soil and Rock samples in the study locations.

1. References [1] United Nations Scientific Committee on Effects of Atomic Radiation. Exposure from Natural Radiation Sources. UNSCEAR Report, New York; 2000.

Primary authors: OHAKWERE-EZE, Michael; Mr NAFIU, Musa

Co-authors: Prof. SINGH, Shiv Kumar; Mr KABIRU, Momoh; Dr SIMON, John

Presenters: Mr NAFIU, Musa; Prof. SINGH, Shiv Kumar; Dr SIMON, John

Session Classification: AfPS Contribution

Track Classification: AfPS