SAIP2013



Contribution ID: 218

Type: Oral Presentation

Extraction of surface impedance from magnetotelluric data

Tuesday, 9 July 2013 16:00 (20 minutes)

Abstract content
 (Max 300 words)

This paper presents the analysis of South African magnetotelluric (MT) data in the time and frequency domain for the purpose of extracting representative values of surface impedance. The surface impedance is used in the derivation of geo-electric fields produced by rapid variations in the geomagnetic field, as occurs during geomagnetic storms. The magnetotelluric method uses the spectra of associated time varying horizontal electric and magnetic fields at the Earth's surface to determine a frequency dependent impedance tensor and an equivalent surface impedance. The theory of operation of MT devices will be presented, as well as typical data obtained from the MT installations in Hermanus, Vaalputs and Middelpos. The various steps in the analysis are aimed at reducing noise and outliers. In the time domain, a Hanning window is used to select data from successive periods during a day, while reducing the end effect (Gibbs' phenomenon) by tapering the series towards the start and ends of each selected time period. The spectral transformation is performed by means of a fast Fourier transformation (FFT). Spectral bands are selected by frequency domain filtering. Typical results and challenges in performing this analysis will be presented.

Apply to be
 considered for a student
 award (Yes / No)?

Yes

Level for award
 (Hons, MSc,
 PhD)?

MSc

Main supervisor (name and email)
and his / her institution

Dr P Cilliers

Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?

Yes

Primary authors: Dr CILLIERS, Pierre (Hermanus Magnetic Observatory); Mr KHANYILE, Sfundo (SANSA Space Science)

Presenter: Mr KHANYILE, Sfundo (SANSA Space Science)

Session Classification: Space Science

Track Classification: Track D2 - Space Science