



Contribution ID: 50

Type: Oral Presentation

Low-Energy Electronic Structure and Fermi Surface of the itinerant metamagnet $\text{Sr}_3\text{Ru}_2\text{O}_7$

Tuesday, 9 July 2013 10:30 (20 minutes)

Abstract content
 (Max 300 words)

$\text{Sr}_3\text{Ru}_2\text{O}_7$ is the second member of the Ruddlesden-Popper series of strongly correlated oxides $\text{Sr}_{n+1}\text{Ru}_n\text{O}_{3n+1}$, where n defines the number of consecutive planes of the ruthenium oxygen (RuO_6) octahedra layers in the unit cell. This compound has recently attracted the attention of the scientific community because of its unusual low-temperature physical properties such as metamagnetism and quantum criticality [1] with evidence of nematic fluid behaviour [2,3]. The early work of Binz et al. based on mean field theory showed that the metamagnetic behaviour observed in $\text{Sr}_3\text{Ru}_2\text{O}_7$ [4] can be understood as a result of the presence of van Hove singularities (vHs) (i.e. sharp peaks observed in the density of states (DOS)) in the proximity of the Fermi energy and associated to Fermi surface (FS) sheets of different symmetry [5]. Therefore, the complete determination of the symmetry and the location of these singularities on the FS are of fundamental importance for the understanding of the quantum critical behaviour of $\text{Sr}_3\text{Ru}_2\text{O}_7$.

In this study we present our recent results on the low-energy electronic structure of $\text{Sr}_3\text{Ru}_2\text{O}_7$, including details on the FS sheets. This investigation was performed via synchrotron-based angle resolved photoemission spectroscopy performed at 1K. Together with the reconstruction of the FS, our results show that - in addition to the flat band close to the X point that reveals a complex DOS with vHs near to the Fermi level (as suggested by Tamai and co-workers [6]) - there is another flat band associated to a FS sheet of a different symmetry around the Γ point that also reveals a complex DOS with vHs near the Fermi level. Moreover, evidence of vHs around the X point has been suggested in the FS of $\text{Sr}_3\text{Ru}_2\text{O}_7$, with the presence of high intensity around this point. Finally, the presence of strong electron-phonon correlations was confirmed by the presence of kinks in band dispersions, which are the evidence of a sensitive coupling between the structural and magnetic properties in $\text{Sr}_3\text{Ru}_2\text{O}_7$.

- [1] S. A. Grigera et al., Science 294, 329 (2001).
- [2] R. A. Borzi et al., Science 315, 214 (2007).
- [3] J. F. Mercure et al., Phys. Rev. Lett. 103, 176401 (2009).
- [4] B. Binz and M. Sigrist, Europhys. Lett. 65, 816 (2004).
- [5] C. Honerkamp, Phys. Rev. B 72, 115103 (2005).
- [6] A. Tamai et al., Phys. Rev. Lett. 101, 026407 (2008).

Apply to be
 considered for a student
 award (Yes / No)?

Yes

Level for award
 (Hons, MSc,
 PhD)?

MSc

**Main supervisor (name and email)
and his / her institution**

Dr. Emanuela Carleschi
Department of Physics,
University of Johannesburg
Auckland Park,2006
ecarleschi@uj.ac.za

**Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?**

Yes

Primary author: Ms SOHANFO NGANKEU, Arlette (Department of Physics University of Johannesburg)

Co-authors: VECCHIONE, Antonio (University of Salerno); DOYLE, Bryan P (Department of Physics, University of Johannesburg); CARLESCHI, Emanuela (Department of Physics, University of Johannesburg); VOBORNIK, Ivana (TASC Trieste); UNNIKRISHNAN, Manju (TASC Trieste); CUOCO, Mario (University of Salerno); FITTI-PALDI, Rosalba (University of Salerno); BORISENKO, Sergey (IFW Dresden); ZABOLOTNYY, Volodymyr (IFW Dresden)

Presenter: Ms SOHANFO NGANKEU, Arlette (Department of Physics University of Johannesburg)

Session Classification: DCMPPM1

Track Classification: Track A - Division for Condensed Matter Physics and Materials