



Contribution ID: 155

Type: Oral Presentation

Research progress in the $H \rightarrow Z\eta^{(*)} \rightarrow \tau^+ \tau^- l^+ l^-$ decay channel

Wednesday, 10 July 2013 09:20 (20 minutes)

Abstract content
 (Max 300 words)

The Higgs to $ZZ^{(*)} \rightarrow l^+ l^- l^+ l^-$ decay channel has good signal to background ratio and can be used to reconstruct the Higgs invariant mass with quite good resolution for an intermediate mass range Higgs ($m_Z < m_H < 2m_Z$). This makes the $ZZ^{(*)}$ channel one of the most relevant channels to study at the LHC, where it has been found that $m_H \approx 125$ GeV. Although the Higgs to $ZZ^{(*)}$ decay channel played an important role in this discovery, the decay leptons considered excluded Z bosons decaying into τ leptons. Reasons for this lie in the hard-to-detect hadronic or leptonic τ decays. A study which includes Z decaying into τ is beneficial to completing the picture of the Higgs search, particularly in the statistically limited vector-boson-fusion like search. Preliminary results of the search for Higgs channel $H \rightarrow ZZ^{(*)} \rightarrow \tau^+ \tau^- l^+ l^-$, ($l = e$ or μ) will be presented and discussed.

Apply to be
 considered for a student
 award (Yes / No)?

No

Level for award
 (Hons, MSc,
 PhD)?

MSc

Main supervisor (name and email)
and his / her institution

Trevor Vickey (Trevor.Vickey@cern.ch) WITS

Would you like to
 submit a short paper
 for the Conference
 Pro- ceedings (Yes / No)?

No

Primary author: Mr HAMITY, Guillermo (Msc Physics (WITS))

Presenter: Mr HAMITY, Guillermo (Msc Physics (WITS))

Session Classification: NPRP

Track Classification: Track B - Nuclear, Particle and Radiation Physics