Contribution ID: **175** Type: **Oral Presentation**

Neutrino Events at IceCube and the Fermi Bubbles

Wednesday, 9 July 2014 14:00 (20 minutes)

Abstract content
   (Max 300 words)
Formatting &
Special chars

The IceCube Collaboration recently announced twenty-eight events were observed with energies above ˜ 30 TeV, more than expected from atmospheric backgrounds. We discuss the detectability of the Fermi Bubbles at IceCube and show that up to 4 – 5 of the 28 events could originate from the Fermi Bubbles (FB). If the observed gamma rays from the FB are created due to the baryonic mechanism, high-energy (> GeV) neutrinos should be emitted as a counterpart. These neutrinos should be detectable as shower or track-like events at a Km3 neutrino detector. For a hard primary cosmic-ray proton spectrum E^{-2.1} and cutoff energy at or above 10 PeV, the Fermi Bubble flux substantially exceeds the atmospheric backgrounds. For a steeper spectrum E^{-2.3} and/or lower cutoff energy, to observe the neutrino flux at high significance, longer running time will be required.

Apply to be
 considered for a student
   award (Yes / No)?

Yes

Level for award
 (Hons, MSc,
   PhD)?

PhD

Main supervisor (name and email)
and his / her institution

Cecilia Lunardini, Cecilia.Lunardini@asu.edu, Arizona State University

Would you like to <br> submit a short paper <br> for the Conference
 Proceedings (Yes / No)?

No

Primary authors: Prof. LUNARDINI, Cecilia (Arizona State University); Dr YANG, Lili (University of Nova Gorica); Prof. RAZZAQUE, Soebur (University of Johannesburg)

Presenter: Dr YANG, Lili (University of Nova Gorica)

Session Classification: Astro

Track Classification: Track D1 - Astrophysics