



Contribution ID: 93

Type: **Oral Presentation**

Modification of the bandoffset in boronitrene

Wednesday, 13 July 2011 12:00 (15 minutes)

We carried out *ab initio* self-consistent calculations to study the modification of the band offsets by polar layers. We considered a double line of C in a monolayer of hexagonal BN, also known as Boronitrene. This effectively introduces a line of dipoles at the interface. We considered BN/C zigzag and armchair configurations. The BN/C zigzag-short is found to be energetically most stable and with the smallest (1.51eV). We show that a two dimensional diode can be constructed from this structure, with changes to the electronic and optical properties of the host system.

**Level (Hons, MSc,
 PhD, other)?**

PhD

**Consider for a student
 award (Yes / No)?**

Yes

**Would you like to
 submit a short paper
 for the Conference
 Proceedings (Yes / No)?**

No

Primary author: Mr OBODO, Kingsley (University of Pretoria, South Africa)

Co-authors: Prof. CHETTY, Nithaya (University of Pretoria, South Africa); Mr ANDREW, Richard C. (University of Pretoria, South Africa)

Presenter: Mr OBODO, Kingsley (University of Pretoria, South Africa)

Session Classification: CMPMS1

Track Classification: Track A - Condensed Matter Physics and Material Science