



Contribution ID: 131

Type: Oral Presentation

## A semiclassical recipe for wobbly limp noodles in partonic soup

Thursday, 6 July 2017 15:00 (20 minutes)

In this talk, I present the main results of our recent paper where we compute the average squared distance travelled by an initially stationary light-flavour off-mass-shell coloured parton in a strongly-coupled thermal plasma using the gauge/string duality. The calculation involves taking an ensemble average of thermal fluctuations induced by Hawking radiation emanating from the black hole horizon in  $\text{AdS}_{d+1}$  - Schwarzschild on top of the leading order worldsheet of our limp noodle string. We argue that our results give us access to the time-dependent transverse momentum squared per unit pathlength picked up by a high momentum quark in a strongly-coupled thermal plasma and present the first time-dependent calculation of the transport coefficient, which is critically important for phenomenology in heavy ion collisions.

**Apply to be considered for a student award (Yes / No)?**

Yes

**Level for award (Hons, MSc, PhD, N/A)?**

MSc

**Main supervisor (name and email) and his / her institution**

William Horowitz wa.horowitz@uct.ac.za

**Would you like to submit a short paper for the Conference Proceedings (Yes / No)?**

Yes

**Primary author:** Mr MOERMAN, Robert (University of Cape Town)

**Co-author:** Dr HOROWITZ, William (University of Cape Town)

**Presenter:** Mr MOERMAN, Robert (University of Cape Town)

**Session Classification:** Theoretical and Computational Physics 1

**Track Classification:** Track G - Theoretical and Computational Physics